

Rethinking Neonatal Vaccination Policies: A Neuroimmune Perspective

Zeinab Paymani¹, MD; Mostafa Nazari¹, MSc; Kayvan Mirnia², MD; Razieh Sangsari², MD; Marzieh Ebrahimi³, PhD; Fatemeh Haghighi⁴, MD

¹Department of Nuclear Medicine, Children Medical Center Hospital, Tehran University of Medical Sciences, Tehran, Iran;

²Department of Neonatology, Children Medical Center Hospital, Tehran University of Medical Sciences, Tehran, Iran;

³Department of Medical Physics, School of Medicine, Iran University of Medical Sciences, Tehran, Iran;

⁴Department of Nuclear Medicine, Children Medical Center Hospital, Tehran University of Medical Sciences, Tehran, Iran

Correspondence:

Mostafa Nazari, MSc;
Children's Medical Center, No. 63,
Gharib Ave., Keshavarz Blvd.,
Postal Code: 1419733151, Tehran, Iran
Tel: +98 21 67412276

Email: msnaa1392@gmail.com

Received: 12 April 2025

Revised: 18 May 2025

Accepted: 19 June 2025

Abstract

Neonatal vaccination is a cornerstone of early-life infectious disease prevention. However, the timing and safety of these interventions require careful consideration. This review explored the neuroimmune implications of early immunization, with a specific focus on the interplay between the developing immune and nervous systems. We examined potential mechanisms through which vaccine-induced immune activation might influence brain development, through epigenetic modifications and sustained cytokine responses, particularly involving interleukin-6 (IL-6). The discussion addressed concerns related to immune overstimulation, regulatory T-cell suppression, and microbiome disruption, considering their potential links to autoimmune and neurodevelopmental disorders. In light of the identified evidence gaps, we advocate for a cautious, individualized vaccination approach guided by the “As Low and Late As Reasonably Achievable” (ALLARA) principle. This strategy aimed to balance robust protection against infectious diseases with the imperative of safeguarding lifelong neurological and immunological health.

Please cite this article as: Paymani Z, Nazari M, Mirnia K, Sangsari R, Ebrahimi M, Haghighi F. Rethinking Neonatal Vaccination Policies: A Neuroimmune Perspective. Iran J Med Sci. 2026;51(2):84-103. doi: 10.30476/ijms.2025.106591.4083.

Keywords • Vaccine • Epigenetics • Autoimmune diseases • Nervous system diseases • Birth dose vaccine

What's Known

- The World Health Organization (WHO) recommends universal administration of birth dose vaccines, including hepatitis B (HBV), oral polio vaccine (OPV), and bacillus Calmette-Guérin (BCG), to prevent early-life infectious diseases. These vaccines are generally considered safe, with reported side effects being predominantly mild and transient, such as low-grade fever, local swelling, or irritability.

What's New

- This review critically examined unexplored dimensions of neonatal vaccination, including immune immaturity, the potential epigenetic implications of early immune activation, and possible autoimmune and neurological consequences. It also addressed challenges of limited immunogenicity and the effectiveness of targeted vaccination strategies. The authors proposed an “As Low and Late As Reasonably Achievable” (ALLARA) based strategy to optimize safety, efficacy, and long-term outcomes of early-life immunization.

Introduction

At birth, both the immune system and the brain of a newborn are functionally immature, possessing only basic capabilities that evolve rapidly during the first 2 years of life.¹ This period is characterized by extensive neurodevelopment, including structural growth, myelination, and connectivity, alongside cognitive, motor, and sensory maturation, all of which influence lifelong behavior.²⁻⁵ Simultaneously, the neonatal immune system is underdeveloped, with reduced functionality in key components, such as monocytes, neutrophils, dendritic cells, natural killer (NK) cells, and T-cells.⁶ Early immunity primarily relies on maternal antibodies transferred via the placenta and breast milk.⁷ Accordingly, studies suggested that immune hyperactivity during fetal and early infancy stages could shape lifelong brain and immune function, potentially increasing disease susceptibility.⁸⁻¹² This concept—that early immune events can have permanent developmental consequences—raises the concern that neonatal exposure to multiple vaccine antigens could alter neuroimmune developmental programs and induce long-term changes in gene expression through epigenetic mechanisms such as DNA methylation or histone modification.

Vaccination represents one of the most effective public health interventions, preventing the spread of infectious diseases and significantly reducing associated morbidity and mortality. Its goal is to elicit a long-lasting, pathogen-specific immune response, while minimizing adverse reactions. It is therefore imperative that vaccine-mediated protection during early life is both safe and efficient. To this end, the World Health Organization (WHO) recommends the administration of specific vaccines, namely the hepatitis B virus (HBV), bacillus Calmette-Guérin (BCG), and oral polio (OPV) vaccine, within the first 24 hours of life. These are referred to as birth-dose vaccines.¹³⁻¹⁵

In this review, we examined neonatal vaccination from a neuroimmune perspective, focusing on its potential epigenetic impacts, safety challenges, efficacy, and current strategies. Special attention was paid to how early immune activation might influence the developing nervous system and long-term health outcomes. This study aimed to provide healthcare professionals and policymakers with evidence-based insights to help guide neonatal vaccination strategies that carefully balance robust immunological protection with neurodevelopmental safety.

Early Life Experience and Lifelong Health

The immune and nervous systems undergo critical, coordinated development during early life, and their interplay significantly affects lifelong health. A growing body of evidence indicate that early life adversity (ELA) could induce long-lasting changes in the immune function, increasing susceptibility to chronic diseases later in life.¹⁶⁻¹⁸ The immune system is integral to normal brain development, behavior, and neural function,¹⁸ and immune dysregulation during sensitive developmental windows may contribute to neurological and psychiatric disorders.¹⁹ Postnatal immune activity has been directly linked to neurological impairments and an increased risk of autoimmune diseases.^{19, 20}

ELA refers to a wide range of adverse exposures—including trauma, stress, infections, and environmental toxins—that can shape immune system development through epigenetic reprogramming.²¹ Such early-life programming may increase the risk of chronic diseases, including cardiovascular, pulmonary, autoimmune, and neurological disorders.²²⁻²⁷ Given that neonatal vaccines are administered during these sensitive developmental periods, it is crucial to investigate their potential epigenetic impacts on the neuroimmune axis (figure 1).

Studies suggested that ELA could lead to heightened innate immune responsiveness and chronic low-grade inflammation, characterized by elevated pro-inflammatory cytokines, such as interleukin-6 (IL-6) and C-reactive protein (CRP).²⁸⁻³⁰ A prominent example is fetal inflammatory response syndrome (FIRS), marked by elevated fetal plasma IL-6 levels in utero, which is associated with an increased risk of neurodevelopmental, psychiatric, autoimmune, cardiovascular, and pulmonary diseases.³¹⁻³³ Similarly, the BCG vaccine—administered at birth in many countries—induces the production of IL-6, interferon gamma (IFN- γ), and tumor necrosis factor alpha (TNF- α), a process known as trained immunity that involves epigenetic reprogramming of innate immunity and monocytes' function.³⁴⁻³⁷ IL-6 levels have been reported to remain elevated for up to a year following BCG vaccination,³⁷ raising concerns about whether this sustained immune activation could represent a form of maladaptive epigenetic reprogramming.

IL-6 is a key cytokine that regulates immune activation, acute phase responses, and tissue repair. However, its chronic elevation is implicated in the pathogenesis of autoimmune disorders, chronic inflammatory diseases, and certain cancers. Consequently, IL-6 blockade has shown therapeutic benefits in experimental models of inflammatory bowel disease, diabetes, multiple sclerosis, asthma, rheumatoid arthritis, and inflammation-related cancers.³⁸



Figure 1: The figure illustrates how early-life exposures, such as stress, trauma, infection, and toxins, can dysregulate the neonatal immune system, contributing to disease in adulthood. It also raises the question of whether neonatal vaccination, as another form of immune-activating factor, could exert similar long-term neuroimmune effects.

Elevated IL-6 also facilitates infiltration of dendritic cells and macrophages into the brain, disrupting neuronal excitability and neurotransmission, which can result in long-term impairments in synaptogenesis and neurogenesis.^{39, 40}

In parallel, the colonization of infant gut microbiota has been widely recognized for its role in brain development and establishing an early life imprint on the immune system.⁴¹ However, alterations to the commensal gut microbiota during this period might increase the risk of inflammatory or allergic diseases later in life. Enterovirus colonization in early infancy, for instance, could restructure the gut microbiome and potentially trigger autoimmunity.⁴² Given this established pathway of virus-induced dysbiosis, an important question arises: could the administration of live attenuated poliovirus via neonatal vaccination contribute to microbiome dysbiosis and an increased risk of autoimmunity?

Newborns possess a uniquely tolerant immunological state, characterized by abundant T regulatory (Treg) cells. These cells are essential for maintaining a balanced and controlled immune system and for preventing inappropriate immune activation. Tregs are key players in controlling inflammation, preventing autoimmunity, and ensuring immune responses are appropriately scaled.⁴³ However, because Tregs can dampen vaccine-induced immunity, adjuvants are often used to suppress Treg activity and enhance immunogenicity.^{44, 45} While this approach supports vaccine efficacy, it may also transiently reduce Treg function in infants. This raises a challenging question: could this early-life reduction in Treg-mediated suppression impact the establishment of lifelong self-tolerance, thereby increasing susceptibility to autoimmunity and future dysregulated immune responses?

Autoimmune and Neurological Adverse Events Following Immunization

A significant body of evidence, including case reports, original articles, reviews, and comparative studies, documented the association between HBV, BCG, and polio vaccines—as well as HBV vaccine adjuvants—and the subsequent development of autoimmune and neurological disorders. However, establishing a causal relationship is challenging. Some of these challenges are discussed in this section.

Vaccine safety surveillance data suggested that most vaccine side effects are usually mild and transient, lasting 1-2 days. However, concerns have been raised regarding the potential for later adverse events.^{46, 47} Furthermore, the durations of pre- and post-license clinical trials are often insufficient for evaluation of long-term

side effects. While many studies indicated that harmful exposures during early life could heighten vulnerability to chronic diseases later in life,^{16-18, 48, 49} scientific literature described several pathways by which vaccines, similar to viruses and other microorganisms, could trigger autoimmune reactions. These include molecular mimicry, cross-reactivity, bystander activation, epitope spreading, and antigen persistence.⁵⁰⁻⁵⁶ For instance, HBV vaccine epitopes were reported in the context of synergistic autoimmune competence.⁵⁷ Additionally, components of the HBV vaccine demonstrated sequence homology and molecular mimicry with human proteins: with the hair follicle protein solute carrier family 45 member 2 (SLC45A2), and with myelin basic protein and myelin oligodendrocyte glycoprotein. These are proposed as plausible biological mechanisms for alopecia areata and multiple sclerosis, respectively.^{58, 59}

We have reviewed studies reporting adverse effects to highlight the potential of these vaccines to contribute to autoimmune and neurological diseases. The cited studies often support a causal link based on a short temporal relationship—typically less than 2 months—between vaccine administration and the appearance of autoimmunity. These adverse effects are not confined to childhood, underscoring the potential for these vaccines to contribute to such conditions across different age groups.

HBV Vaccine Adverse Events

Multiple case reports and case series highlighted a connection between the HBV vaccine and various autoimmune and neurological diseases, including arthritis/polyarthralgia, lupus erythematosus, multiple sclerosis, optic neuritis, vasculitis, alopecia areata, erythema nodosum, polyarteritis nodosa (PAN), thrombocytopenic purpura, evans syndrome, Guillain-Barré Syndrome (GBS), glomerulonephritis, uveitis, polymyositis, dermatomyositis, Takayasu's arteritis, Hashimoto's thyroiditis, Graves' disease, childhood bullous pemphigoid, chronic fatigue syndrome, cutaneous pseudo lymphoma, vitiligo, lichen planus. A comprehensive list is presented in table 1.

In addition to the side effects associated with HBV vaccine epitopes, HBV vaccines contain aluminum adjuvants as boosters of immune response. The adjuvants are substances added to vaccines to enhance the immunogenicity of the vaccine antigens.⁹⁷⁻⁹⁹ Studies indicated that aluminum-based adjuvants in the HBV vaccine are associated with neuropsychiatric symptoms, fatigue, mucocutaneous, musculoskeletal, and gastrointestinal complaints.

Table 1: Autoimmune and neurological disorders post hepatitis B vaccine

Author, year, reference	Events	Type of study
Geier et al., 2005 ⁵⁷	Multiple sclerosis, optic neuritis, vasculitis, arthritis, alopecia, lupus erythematosus, rheumatoid arthritis	Case control
Geier M et al., 2003 ⁶⁰	Erythema nodosum, lichen planus, polyarteritis nodosa, Reiter syndrome, thrombocytopenic purpura, Evans syndrome, acute posterior multifocal placoid pigment epitheliopathy, optic neuritis, transverse myelitis, central nervous system demyelination, cerebellar ataxia, multiple sclerosis, chronic fatigue syndrome	Review
Maubec et al., 2005 ⁶¹	Cutaneous pseudolymphoma, vitiligo, chronic fatigue syndrome	Case series
Júnior et al., 2020 ⁶²	Graves' Disease, rheumatoid arthritis (RA), psoriasis, lupus, Hashimoto's thyroiditis (HT), vitiligo	Case control
Oscar-Danilo et al., 2009 ⁶³	Chronic fatigue syndrome	Review
Mikaeloff et al., 2009 ⁶⁴	Multiple sclerosis	Case control
Herroelen et al., 1991 ⁶⁵	Multiple sclerosis	Case reports
Tourbah et al., 1999 ⁶⁶	Central nervous system demyelination	Case series
Agmon-Levin et al., 2014 ⁶⁷	Chronic fatigue syndrome, fibromyalgia	Case control
Nancy et al., 2008 ⁶⁸	Chronic fatigue syndrome	Case reports
Richardson et al., 2018 ⁶⁹	Alopecia areata	Case reports
Choffray et al., 2007 ⁷⁰	Lupus panniculitis	Case reports
Luhadia et al., 2022 ⁷¹	Multiple sclerosis	Case reports
Case Series et al., 2002 ⁷²	Lichen planus	Case series
de la Fuente et al., 2013 ⁷³	Childhood bullous pemphigoid	Case series
Erbagci et al., 2002 ⁷⁴	Childhood bullous pemphigoid	Case reports
Berkun et al., 2005 ⁷⁵	Pemphigus	Case reports
Vital et al., 2002 ⁷⁶	Inflammatory neuropathy	Case reports
De Carvalho et al., 2008 ⁷⁷	Systemic polyarteritis nodosa	Review
Maillefert et al., 1997 ⁷⁸	Polyarthralgia	Case reports
Zaas et al, 2001 ⁷⁹	Takayasu's arteritis	Case reports
Agmon-Levin et al., 2009 ⁸⁰	Systemic lupus erythematosus	Case series
Bogdanos et al., 2009 ⁵⁸	Multiple sclerosis	Case control
Altman et al., 2008 ⁸¹	Dermatomyositis	Case reports
Geier et al., 2004 ⁸²	Arthritis, rheumatoid arthritis, myelitis, optic neuritis, multiple sclerosis, Guillain-Barré syndrome, glomerulonephritis, thrombocytopenia, systemic lupus erythematosus	Case series
Geier et al., 2002 ⁸³	Arthralgia, arthrosis, arthritis, thrombocytopenia, hepatitis, erythema, Guillain-Barré Syndrome, myelitis, vasculitis	VAERS
Pennesi et al., 2002 ⁸⁴	Glomerulonephritis	Case reports
Poierriez J et al., 2004 ⁸⁵	Transverse myelitis, neurolupus	Case reports
Schattner et al., 2005 ⁸⁶	Rheumatoid arthritis, reactive arthritis, vasculitis, encephalitis, neuropathy, thrombocytopenia	Review
Cohen et al., 1996 ⁵⁴	Erythema nodosum, immune thrombocytopenia, myasthenia gravis, uveitis, Reiter's syndrome, arthritis, systemic lupus erythematosus, central nervous system demyelination, anti-DNA antibodies emergence, Evans' syndrome	Review
Ramirez Rivera et al., 2003 ⁸⁷	Polymyositis	Case report
Agmon-Levin, 2009 ⁸⁸	Transverse myelitis with a short interval <2 months	Multianalysis
Maillefert et al., 1999 ⁸⁹	Rheumatoid arthritis, systemic lupus erythematosus, polyarthralgia, myalgia, vasculitis, miscellaneous with mixed presentations	Original article
Ronch et al., 1998 ⁹⁰	Immune thrombocytopenic in infants within 1 month	Case series
Neau et al., 1998 ⁹¹	Immune thrombocytopenic in children	Case series
Berkun et al., 2005 ⁷⁵	Pemphigus	Case reports
Chave et al., 2003 ⁹²	Henoch shonlein purpura	Case report
Khamaisi et al., 2004 ⁹³	Guillain-Barré syndrome	Case reports
Girard et al., 2004 ⁹⁴	Multiple sclerosis, chronic fatigue syndrome	Review
Wise et al., 1997 ⁹⁵	Alopecia	Case Series
Avci et al., 2013 ⁹⁶	Hemolytic uremic syndrome	Case report

VAERS: vaccine adverse event reporting system

They have also been linked to autoimmune/inflammatory syndrome induced by adjuvants (ASIA) syndrome, autism spectrum disorder, sarcoidosis, Sjogren's syndrome, elevated

titers of autoantibodies, and undifferentiated connective tissue diseases, as summarized in table 2. Aluminum compounds can persist in the human body for many years post-vaccination.¹⁰⁰

Table 2: Autoimmune and neurological disorders linked to aluminum adjuvants

Author, year, reference	Events	Type of study
Shoenfeld et al., 2011 ¹⁰²	Macrophagic myofasciitis syndrome (MMF) Gulf War Syndrome	Review
Zafirir et al., 2012 ¹⁰³	Neuro-psychiatric symptoms, fatigue, mucocutaneous, musculoskeletal, and gastrointestinal complaints. Elevated titers of autoantibodies	Original
Boretti et al., 2021 ¹⁰¹	Autism	Review
Shaw, et al., 2013 ¹⁰⁴	Autoimmune/inflammatory syndrome, autism spectrum disorders	Review
Graham Ewing, 2009 ¹⁰⁵	Autism	Review
Tomljenovic et al., 2011 ¹⁰⁶	Autoimmunity, long-term brain inflammation, and associated neurological complications	Review
Borba, 2020 ¹⁰⁷	Sarcoidosis, Sjögren's syndrome, undifferentiated connective tissue disease	Review

Table 3: Autoimmune and neurological disorders linked to bacillus Calmette-Guérin (BCG) and intravesical BCG (iBCG) vaccine

Author, year, reference	Events	Type of study
Jain et al., 2022 ¹⁰⁹	Phlyctenular conjunctivitis, scleritis, sclerokeratitis, retinal periphlebitis, iridocyclitis, choroiditis, uveitis, keratitis	Review
Khalili et al., 2021 ¹¹⁰	Chronic granulomatous disease	Case reports
Dahl et al., 2020 ¹¹¹	Increased risk of hip fracture	Cohort
Tsujioka et al., 2022 ¹¹²	Osteomyelitis	Cohort
Wang et al., 2022 ¹¹³	Lymphadenitis	Cohort
Sellami et al., 2018 ¹¹⁴	Disseminated bacillus Calmette-Guérin	Cohort
Modrzejewska et al., 2006 ¹¹⁵	Detachment of retina	Case reports
Salmon et al., 2019 ¹¹⁶	Hodgkin's lymphoma (HL)	Cohort
Shoenfeld et al., 2001 ¹¹⁷	Reiter's syndrome	Case reports
Schuchmann et al., 2001 ¹¹⁸	Juvenile chronic arthritis	Case reports
Anis et al., 2023 ¹¹⁹	Reactive arthritis, psoriasis, Myasthenia gravis, ocular manifestations	Review
Sharan et al., 2005 ¹²⁰	Autoimmune retinopathy	Case reports
Nakagawa et al., 2018 ¹²¹	Reiter's syndrome	Case series
Genereau et al., 1996 ¹²²	Polymyalgia rheumatica	Case reports
Thepot et al., 1995 ¹²³	Acute polyarthritis	Case reports
Granel et al., 2004 ¹²⁴	Cryoglobulinemic vasculitis	Case reports
Tsuchiya et al., 2021 ¹²⁵	Intestinal ulcers	Case reports
Parent et al., 2018 ¹²⁶	Primary angiitis of the central nervous system (PACNS)	Review
Beisland et al., 2004 ¹²⁷	Vitiligo	Case reports
Shoenfeld, 2001 ¹¹⁷	Inflammatory arthritis	Case series
Bernini, et al., 2015 ⁴⁶	Kawasaki, arthritis	Review
Vittori, et al., 1996 ¹²⁸	Tuberculosis lupus	Case reports
Izumi, 1982 ¹²⁹	Lupus vulgaris	Case reports
Noishiki et al., 2023 ¹³⁰	Keloid	Review
Abid et al., 2021 ¹³¹	Acute hepatitis with granulomatous steatohepatitis and portal fibrosis	Case reports
Yamazaki-Nakashimada et al., 2019 ¹³²	Endophthalmitis, uycotic aneurysms, Takayasu arteritis, Kawasaki disease, Reiter syndrome, and Guillain-Barré syndrome	Review
Sumida, et al., 2003 ¹³³	Autoimmune pancreatitis	Case report
Foucard, et al., 1971 ¹³⁴	Osteoarthritis	Case series

Moreover, aluminum has demonstrated a detrimental impact on the blood-brain barrier (BBB) and is connected to microglia-triggered pro-inflammatory cytokine release. Due to its high reactivity, the aluminum ion (Al³⁺) can interfere with several biological functions in the developing brain, including enzymatic activities of key metabolic pathways. In the context of infancy, a significant correlation has been reported between pediatric vaccines containing aluminum adjuvants and the incidence of autism spectrum disorders. Infants receiving these vaccines have a notably higher incidence of

autism spectrum disorder (ASD), suggesting a potential association between these vaccine components and developmental sequelae.¹⁰¹

BCG Vaccine Adverse Events

Previous studies reported autoimmune and neurological disorders following BCG vaccination. A portion of the evidence regarding BCG adverse events is derived from studies utilizing intravesical BCG (iBCG) for cancer immunotherapy. Reported side effects include juvenile idiopathic arthritis (JIA), juvenile dermatomyositis, Takayasu arteritis, autoimmune pancreatitis, GBS, optic neuritis,

meningitis, vasculitis, psoriasis, endophthalmitis, uveitis, autoimmune retinopathies, Hodgkin's lymphoma, lymphadenitis, osteomyelitis, osteitis, and disseminated disease (BCGosis), as illustrated in table 3.

Besides, BCG vaccination has presented a significantly high rate of complications in patients with severe combined immunodeficiency (SCID), leading to substantial morbidity and mortality. An analysis of BCG-vaccinated patients with SCID from 28 centers across 17 countries revealed that early vaccination (≤ 1 month) was associated with a higher prevalence of BCG-related complications and death.¹⁰⁸

OPV Vaccine Adverse Events

Although adverse events related to the oral polio vaccine (OPV) are generally considered rare, cases of autoimmune effects following colonization of the gut by the attenuated polioviruses, a serotype of enterovirus C within the *picornaviridae* family, have been reported. Autoimmune events associated with OPV include multiple sclerosis, childhood acute disseminated encephalomyelitis, vaccine-associated paralytic poliomyelitis (VAPP), acute flaccid paralysis, immune thrombocytopenia (ITP), Gianotti-Crosti syndrome (GCS), transverse myelitis, ulcerative colitis (UC), and Crohn's disease (CD), as detailed in table 4.

Challenges of Discovering Adverse Events Post-Vaccination

The WHO provides a comprehensive guideline for assessing causality in adverse events following immunization (AEFI), emphasizing well-defined clinical documentation, a temporal association with vaccination, biological plausibility, and the exclusion of alternative causes.¹⁴⁴ This framework prioritizes identifying strong alternative explanations—such as genetic or pre-existing conditions—before attributing

events to vaccines. While this method ensures scientific rigor, it may overlook the complex, multifactorial nature of autoimmune and neurologic diseases, particularly when vaccines act as contributing factors rather than single causes. Moreover, although biological plausibility and timing are central to the WHO's approach, vaccines can affect the developing immune and nervous systems—especially in early life—with clinical manifestations potentially appearing years later. This underscores the critical need for long-term monitoring. Finally, emerging research is essential to reveal pathophysiological mechanisms that are not yet fully understood.

The WHO guideline also depends heavily on existing literature to assess and often exclude causal links. However, the current literature is often derived from passively gathered data, such as that in the vaccine adverse event reporting system (VAERS), which has inherent methodological limitations. These include a lack of systematic follow-up, significant underreporting, and a failure to capture delayed onset conditions, particularly autoimmune or chronic diseases that manifest long after vaccination. Furthermore, there is no clear guidance on appropriate time windows for monitoring such delayed-onset diseases. It is therefore inappropriate to assume that autoimmune side effects are rare in the absence of robust and comprehensive documentation. Overlapping symptoms and intensive infant vaccination schedules further complicate the identification of specific causal relationships.

To enhance causality assessment, a robust, multidisciplinary approach involving epidemiologists, clinicians, immunologists, and basic scientists is essential. Such collaboration can help elucidate the nuanced relationships between vaccination and long-term immunological or neurological outcomes, thereby informing evidence-based public health strategies.

Table 4: Autoimmune and neurological disorders linked to OPV vaccine

Author, year, reference	Events	Type of study
Zawar et al., 2017 ¹³⁵	Gianotti-Crosti syndrome	Cohort
Gao et al., 2021 ¹³⁶	Immune thrombocytopenic purpura (ITP), vaccine-associated paralytic poliomyelitis (VAPP)	Cohort
Akbayram et al., 2015 ¹³⁷	Immune thrombocytopenic purpura (ITP)	Cohort
Elkhatay et al., 2020 ¹³⁸	Childhood acute disseminated encephalomyelitis	Case series
Kelly et al., 2006 ¹³⁹	Transverse myelitis	Review
Hughes et al., 2020 ¹⁴⁰	Central nervous system demyelination	Cohort
Agmon-Levin et al., 2009 ⁸⁸	Transverse myelitis with a short interval <2 months	Multi-analysis
Chambrun et al., 2015 ¹⁴¹	Crohn's, ulcerative colitis	Review
Plat et al., 2014 ¹⁴²	Paralytic poliomyelitis (VAPP)	Review
Ami Schattner et al., 2005 ⁸⁶	Neurologic (encephalitis, Guillain-Barré syndrome) Rheumatic (acute arthritis) Hematologic (thrombocytopenia, immune hemolytic anemia)	Review
Dhiman et al., 2018 ¹⁴³	Acute flaccid paralysis	Original

Another avenue for exploring causal relationships and potential side effects is to compare health outcomes between vaccinated versus unvaccinated populations. Although such studies are often not feasible due to high vaccination coverage in many countries, a limited number of observational surveys exist (table 5). These studies reported higher prevalence of developmental delays, severe allergies, attention-deficit hyperactivity disorder (ADHD), autism spectrum disorders, and even infections (e.g., pneumonia and otitis media) in vaccinated cohorts. Future studies with a two-group design are required to provide further insights.

Efficacy of Neonatal Vaccines

The neonatal immune system is characterized by a state of immune tolerance. At this stage of life, all circulating antibodies are of maternal origin, providing passive immunity until the newborn's own antibody production becomes robust around 3 months after birth.^{7, 148-150} These maternal antibodies are highly effective against most infections.⁷ Due to the inherent immaturity of the immune system, neonatal innate immunity relies on distinctive mechanisms. In response to pathogens, the innate immune system serves as the first line of defense. However, neonatal monocytes and dendritic cells (DCs) produce less TNF, IL-12, and IFN- γ , while increasing the production of IL-6, IL-10, and IL-23.^{151, 152} Neutrophils exhibit quantitative and qualitative differences compared to those in older children.¹⁵³ NK cells display diminished cytotoxic capability and impaired release of

destructive substances against infected cells.¹⁵⁴ Furthermore, neonatal NK cells release less IFN- γ , and their adhesion is compromised due to decreased expression of specific adhesion molecules.¹⁵⁵ This immunological bias renders newborns prone to low inflammatory responses and impairs their responses to many vaccines.¹⁵⁶ In this section, we examine the efficacy of BCG, HBV, and OPV vaccines during the neonatal period. The results are summarized in table 6.

HBV Vaccine Efficacy

Hepatitis B surface antibodies (anti-HBs) are produced by the immune system in response to the hepatitis B surface antigen and serve as a marker for immunity.¹⁶⁸⁻¹⁷⁰ Vaccine efficacy, an anti-HBs level >10 IU/L after vaccination, provides complete protection against acute and chronic hepatitis B.^{15, 171} The HBV vaccine is highly effective in infants, with over 95% of healthy recipients developing seroprotective anti-HBs levels within 1 month after the final dose.^{172, 173} However, few studies have evaluated the seroprotective rate of the HBV vaccine specifically in the neonatal period. These studies reported seroconversion rates of only 18% to 50%, 1 month after receiving the birth dose, without accounting for the potential interference of maternal antibodies (table 6).^{157, 174} Moreover, several studies indicated that individuals with a low antibody response exhibited reduced T-cell proliferation and cytokine production.^{158, 175-177} Research also showed that the immune response was enhanced when the first vaccination dose was administered at 2 months of age.^{174, 178-180}

Table 5: Comparative studies between vaccinated versus unvaccinated children

Author, year, reference	Vaccinated vs. unvaccinated	Type of study
Hooker et al., 2020 ¹⁴⁵	Developmental delays (OR=2.18), asthma (OR=4.49), and otitis media (OR=2.13).	Comparative study
Hooker et al., 2021 ¹⁴⁶	Severe allergies (OR=4.31), autism (OR=5.03), gastrointestinal disorders (OR=13.8), asthma (OR=17.6), ADHD (OR=20.8), Chronic ear infections (OR=27.8)	Comparative study
Mawson et al., 2017 ¹⁴⁷	Neurodevelopmental delay, pneumonia, otitis media, allergies	Comparative study

Table 6: Efficacy of hepatitis B, bacillus Calmette-Guérin, and oral polio vaccine in one month of life

Author, year, reference	Vaccine	Efficacy (%)	Time measurement
Soulié et al., 1991 ¹⁵⁷	HBV	50	4 weeks
Strandmark et al., 2022 ¹⁵⁸	HBV	18	4 weeks
Martinez et al., 2022 ¹⁵⁹	BCG	42	<3 years
Roy et al., 2014 ¹⁶⁰	BCG	32	<5 years
Waggie et al., 2011 ¹⁶¹	OPV	41	4 weeks
Sutter et al., 2010 ¹⁶²	OPV	15	4 weeks
El-Sayed et al., 2008 ¹⁶³	OPV	37	4 weeks
Bhaskaram et al., 1997 ¹⁶⁴	OPV	4	6 weeks
Jain et al., 1997 ¹⁶⁵	OPV	13	6 weeks
Khare et al., 1993 ¹⁶⁶	OPV	38	6 weeks
Dong et al., 1986 ¹⁶⁷	OPV	37	4 weeks

HBV: Hepatitis B; BCG: Bacillus Calmette-Guérin; OPV: Oral polio vaccine

This improvement could be attributed to the maturation of the infant's immune system.¹⁸¹⁻¹⁸³ A large review in Africa demonstrated that children born to HBsAg-negative mothers, the risk of infection remained minimal even when vaccination began at 2 months, suggesting no clear additional benefits from the HBV birth dose.¹⁸⁴

BCG Vaccine Efficacy

The BCG vaccine has stood as the exclusive vaccine against tuberculosis (TB) for decades.¹⁴⁶ We did not identify any studies specifically reporting the efficacy of BCG vaccination in the early neonatal period. However, a study reported 42% efficacy in children under 5 years of age,¹⁶⁰ while another showed 32% efficacy in children under 3 against all forms of tuberculosis.¹⁵⁹

A limited number of studies explored the immune responses following neonatal BCG vaccination. These investigations have identified CD4-positive (CD4+) and CD8-positive (CD8+) T lymphocytes as the predominant responding cell populations.^{36, 185} The CD4+ T-cells notably upregulate IFN- γ , TNF- α , IL-2, and IL-6, whereas CD8+ T-cells demonstrate minimal to undetectable production of IFN- γ , TNF- α , and IL-2.^{35, 36, 186} However, the reliability of BCG-specific CD4+ and CD8+ T-cell cytokine expression as a correlate of protection against childhood TB has been questioned.^{35, 187} The limited efficacy of vaccination confirms this theory.¹⁸⁸

Furthermore, investigations revealed that Th1 immune responses become detectable approximately 2-3 months post-vaccination.^{187, 189} Studies have also indicated that immunogenicity is enhanced when BCG administration is postponed until 10 weeks of age.¹⁸⁶

OPV Vaccine Efficacy

Multiple studies have evaluated the effectiveness of the birth dose of the live attenuated oral poliovirus vaccine (OPV). The observed seroconversion rates revealed a range of responses across the different poliovirus serotypes. For type 1, seroconversion rates spanned from 6% to 42% (mean=28%), for type 2, the rates ranged from 2% to 63% (mean=36%), and for type 3, the seroconversion rates varied between 2% and 35% (mean=16%).¹⁶¹⁻¹⁶⁷ This considerable variability underscored the limited and unpredictable immune response triggered by the OPV birth dose in the neonatal period. A more robust immune response is observed with increasing age, highlighting the critical importance of timing and subsequent booster doses for achieving

reliable protection against poliovirus infection.

Current Neonatal Vaccination Strategies

The Expanded Program on Immunization (EPI) was established by the WHO in 1974 with the initial goal of protecting children against six major diseases: tuberculosis, polio, diphtheria, tetanus, pertussis, and measles. The program has since expanded its scope to include additional vaccines and immunization coverage goals. A key component of the EPI is the "birth dose"—the administration of a vaccine shortly after birth to provide early protection against diseases that pose an immediate risk to newborns. This strategy is critical for preventing mother-to-child or early environmental transmission of specific infections.

According to WHO guidelines, the birth dose includes specific vaccines to be given within the first 24 hours of life: the BCG vaccine, the zero dose of OPV, and the HBV vaccine.¹³⁻¹⁵ However, the specific vaccines included and their exact timing can vary based on national health policies and local disease prevalence.

Globally, two primary vaccination strategies are typically implemented for newborns: the 'general recommendation' strategy and the 'recommendation to at-risk groups' strategy. A general recommendation strategy for neonates involves compulsory administration of essential vaccines to all newborns to establish early protection against preventable diseases. In contrast, the 'recommendation for at-risk groups', also known as the selective or targeted strategy, focuses on identifying and prioritizing neonates who face a higher likelihood of exposure to specific infectious diseases or an increased risk of complications due to underlying health conditions or environmental factors.

Nations worldwide implement distinct childhood immunization strategies to protect infants from infectious diseases. While some countries adopt a universal approach, vaccinating all newborns irrespective of their risk factors, others employ a targeted strategy, focusing on specific at-risk groups. Several countries, including Austria, Belgium, the Czech Republic, Denmark, Germany, Iceland, Italy, the Netherlands, Slovakia, and Spain, have discontinued universal BCG vaccination, removing it from their routine schedules. Others, such as Cyprus, Finland, France, Norway, Slovenia, Sweden, and the United Kingdom, currently recommend it exclusively for specific at-risk categories of children. This approach notably includes those with parents from high-TB-prevalence countries or with a family history of TB. The rationale for this targeted approach




Figure 2: This roadmap illustrates how critical assessment of four key factors—epigenetic impacts, safety challenges, efficacy, and current strategies—highlights the potential influence of early-life immune activation on long-term health. This integrated analysis supports a shift toward more cautious and evidence-based vaccination approaches.

in low-endemic countries is based on the low infection risk, a high number needed to vaccinate (NNV), i.e., the number of healthy individuals who must be vaccinated to prevent one case of TB, and a high rate of adverse events per prevented TB case.

A similar pattern is seen with the HBV birth dose. Some countries, including Austria, Cameroon, Finland, Germany, Greece, Hungary, Iceland, and Ireland, have not included it in their routine immunization schedules. In contrast, several countries, such as Canada, Belgium, Czechia, Denmark, Estonia, France, Italy, Japan, Latvia, Luxembourg, the Netherlands, New Zealand, Norway, San Marino, Slovakia, Spain, Sweden, Switzerland, and the United Kingdom, recommend HBV birth dose for at-risk groups. This selective recommendation reflects a targeted public health approach, likely considering factors, such as regional prevalence, disease severity, and available resources. Furthermore, OPV is administered in many countries from 2 months of age. Such diverse approaches highlight the adaptability of immunization strategies to individual country requirements and health priorities.

Neonate Vaccination: Future Directions

Neonatal vaccination plays a critical role in preventing early-life infectious diseases, yet it presents unique challenges due to the immaturity of the infant’s immune system. The limited immunogenic responses in neonates can reduce vaccine efficacy and raise concerns

regarding the long-term impact of early immune activation. As the immune system matures with age, a more robust and balanced response can be achieved, supporting the consideration of delayed or staged immunization schedules.

Recent concerns have focused on the potential neuroimmune effects of immune stimulation during sensitive developmental windows. Emerging evidence suggested that early-life immune activation might influence epigenetic programming and increase the risk of autoimmune and neurodevelopmental disorders later in life. In response, several countries have begun adopting selective neonatal vaccination strategies, prioritizing high-risk infants and integrating early screening programs. This shift offers valuable opportunities to evaluate long-term outcomes between vaccinated and unvaccinated populations and to tailor immunization strategies more precisely. Figure 2 conceptually outlines these four key considerations—epigenetic impacts, safety challenges, efficacy, and current strategies—guiding a more cautious and evidence-based reevaluation of neonatal vaccination policy.

Conclusion

Vaccination remains a cornerstone of pediatric health. However, its application in the neonatal period requires careful consideration. The balance between providing early protection against infectious diseases and the potential risks of overstimulating the developing immune system

must be guided by rigorous scientific evidence. A growing body of research highlight that neonatal immune responses differ significantly from those in older children, necessitating an individualized, developmentally informed approach to vaccine scheduling and administration.

As technologies such as artificial intelligence and precision medicine advance, they offer unprecedented opportunities to design personalized vaccine schedules based on an infant's genetic predispositions and environmental risk factors. Adopting a flexible, risk-based framework—aligned with the ALLARA principle—can help optimize both safety and efficacy of early-life immunization. Ultimately, neonatal vaccination policies must prioritize long-term neurological and immunological well-being, ensuring that each administered vaccine is both necessary and appropriate for the individual infant.

Acknowledgment

We would like to express our gratitude to all those who have encouraged a critical and balanced perspective on vaccination, reminding us to view vaccines—resembling any medical intervention—as procedures that require careful consideration of both risks and benefits, rather than assuming them to be inherently risk-free.

Authors' Contribution

Z.P: Contributed to study conception and design, data analysis, manuscript revision; M.N: Responsible for data acquisition, data interpretation, drafting major sections of the manuscript; K.M: Supervised the project, contributed to data interpretation, critically revised the manuscript; R.S: Assisted in study design, coordinated clinical data collection, contributed to drafting; M.E: Performed data analysis, ensured data quality, participated in manuscript revision; F.H: Conducted the literature review, contributed to data interpretation, assisted in manuscript preparation. All authors have read and approved the final manuscript and agree to be accountable for all aspects of the work in ensuring that questions related to the accuracy or integrity of any part of the work are appropriately investigated and resolved.

Declaration of AI

We declare that AI tools (Chat gpt-4) were used solely to assist with language editing, grammar correction, and improvement of clarity in the manuscript. AI tools were not used to generate

original scientific content, analyze data, interpret results, or draw conclusions. All study design, data analysis, interpretation of results, and scientific judgments were performed by the authors. The authors take full responsibility for the accuracy, originality, and integrity of the work and confirm that the manuscript complies with the journal's ethical and authorship guidelines.

Conflict of Interest: None declared.

References

- 1 Bateson P, Barker D, Clutton-Brock T, Deb D, D'Udine B, Foley RA, et al. Developmental plasticity and human health. *Nature*. 2004;430:419-21. doi: 10.1038/nature02725. PubMed PMID: 15269759.
- 2 Dekaban AS. Changes in brain weights during the span of human life: relation of brain weights to body heights and body weights. *Ann Neurol*. 1978;4:345-56. doi: 10.1002/ana.410040410. PubMed PMID: 727739.
- 3 Richards JE, Conte S. Brain Development in Infants: Structure and Experience. In: Lockman JJ, Tamis-LeMonda CS, editors. *The Cambridge Handbook of Infant Development: Brain, Behavior, and Cultural Context*. Cambridge Handbooks in Psychology. Cambridge: Cambridge University Press; 2020. p. 94-127.
- 4 Gilmore JH, Knickmeyer RC, Gao W. Imaging structural and functional brain development in early childhood. *Nat Rev Neurosci*. 2018;19:123-37. doi: 10.1038/nrn.2018.1. PubMed PMID: 29449712; PubMed Central PMCID: PMC5987539.
- 5 Tierney AL, Nelson CA, 3rd. Brain Development and the Role of Experience in the Early Years. *Zero Three*. 2009;30:9-13. PubMed PMID: 23894221; PubMed Central PMCID: PMC3722610.
- 6 Georgountzou A, Papadopoulos NG. Postnatal Innate Immune Development: From Birth to Adulthood. *Front Immunol*. 2017;8:957. doi: 10.3389/fimmu.2017.00957. PubMed PMID: 28848557; PubMed Central PMCID: PMC554489.
- 7 Cinicola B, Conti MG, Terrin G, Sgrulletti M, Elfeky R, Carsetti R, et al. The Protective Role of Maternal Immunization in Early Life. *Front Pediatr*. 2021;9:638871. doi: 10.3389/fped.2021.638871. PubMed PMID: 33996688; PubMed Central PMCID: PMC8113393.
- 8 Danese A, S JL. Psychoneuroimmunology of Early-Life Stress: The Hidden Wounds

- of Childhood Trauma? Neuropsychopharmacology. 2017;42:99-114. doi: 10.1038/npp.2016.198. PubMed PMID: 27629365; PubMed Central PMCID: PMC5143500.
- 9 Eriksson M, Raikkonen K, Eriksson JG. Early life stress and later health outcomes--findings from the Helsinki Birth Cohort Study. *Am J Hum Biol.* 2014;26:111-6. doi: 10.1002/ajhb.22502. PubMed PMID: 24375506.
 - 10 Spitzer C, Bouchain M, Winkler LY, Wingenfeld K, Gold SM, Grabe HJ, et al. Childhood trauma in multiple sclerosis: a case-control study. *Psychosom Med.* 2012;74:312-8. doi: 10.1097/PSY.0b013e31824c2013. PubMed PMID: 22408134.
 - 11 Laforest-Lapointe I, Arrieta MC. Patterns of Early-Life Gut Microbial Colonization during Human Immune Development: An Ecological Perspective. *Front Immunol.* 2017;8:788. doi: 10.3389/fimmu.2017.00788. PubMed PMID: 28740492; PubMed Central PMCID: PMC5502328.
 - 12 Hata M, Hata M, Andriessen EM, Juneau R, Pilon F, Crespo-Garcia S, et al. Early-life peripheral infections reprogram retinal microglia and aggravate neovascular age-related macular degeneration in later life. *J Clin Invest.* 2023;133. doi: 10.1172/JCI159757. PubMed PMID: 36787231; PubMed Central PMCID: PMC5927938.
 - 13 BCG vaccines: WHO position paper - February 2018. *Wkly Epidemiol Rec.* 2018;93:73-96. PubMed PMID: 29474026.
 - 14 Polio vaccines: WHO position paper - March, 2016. *Wkly Epidemiol Rec.* 2016;91:145-68. PubMed PMID: 27039410.
 - 15 Hepatitis B vaccines: WHO position paper - July 2017. *Wkly Epidemiol Rec.* 2017;92:369-92. PubMed PMID: 28685564.
 - 16 Nobile S, Di Sipio Morgia C, Vento G. Perinatal Origins of Adult Disease and Opportunities for Health Promotion: A Narrative Review. *J Pers Med.* 2022;12. doi: 10.3390/jpm12020157. PubMed PMID: 35207646; PubMed Central PMCID: PMC8877993.
 - 17 Bilbo SD, Schwarz JM. Early-life programming of later-life brain and behavior: a critical role for the immune system. *Front Behav Neurosci.* 2009;3:14. doi: 10.3389/neuro.08.014.2009. PubMed PMID: 19738918; PubMed Central PMCID: PMC2737431.
 - 18 Brydges NM, Reddaway J. Neuroimmunological effects of early life experiences. *Brain Neurosci Adv.* 2020;4:2398212820953706. doi: 10.1177/2398212820953706. PubMed PMID: 33015371; PubMed Central PMCID: PMC7513403.
 - 19 Kim J, Erice C, Rohlwick UK, Tucker EW. Infections in the Developing Brain: The Role of the Neuro-Immune Axis. *Front Neurol.* 2022;13:805786. doi: 10.3389/fneur.2022.805786. PubMed PMID: 35250814; PubMed Central PMCID: PMC8891478.
 - 20 Robaina Castellanos GR, Riesgo Rodriguez Sde L. Neonatal sepsis and neurodevelopment in very low birth weight infants in Matanzas, Cuba 2006-2010: a prospective cohort study. *Medwave.* 2016;16:e6422. doi: 10.5867/medwave.2016.03.6422. PubMed PMID: 27097362.
 - 21 Grova N, Schroeder H, Olivier JL, Turner JD. Epigenetic and Neurological Impairments Associated with Early Life Exposure to Persistent Organic Pollutants. *Int J Genomics.* 2019;2019:2085496. doi: 10.1155/2019/2085496. PubMed PMID: 30733955; PubMed Central PMCID: PMC6348822.
 - 22 Rocca WA, Gazzuola Rocca L, Smith CY, Esterov D, Kapoor E, St Sauver JL, et al. Adverse childhood experiences increase the long-term accumulation of morbidity in women. *Commun Med (Lond).* 2025;5:287. doi: 10.1038/s43856-025-00961-0. PubMed PMID: 40640470; PubMed Central PMCID: PMC6348822.
 - 23 Wu Y, Xu R, Gasevic D, Yang Z, Yu P, Wen B, et al. UK Biobank data demonstrate long-term exposure to floods is a risk factor for incident dementia. *Commun Med (Lond).* 2025;5:71. doi: 10.1038/s43856-025-00771-4. PubMed PMID: 40065151; PubMed Central PMCID: PMC6348822.
 - 24 Alley J, Gassen J, Slavich GM. The effects of childhood adversity on twenty-five disease biomarkers and twenty health conditions in adulthood: Differences by sex and stressor type. *Brain Behav Immun.* 2025;123:164-76. doi: 10.1016/j.bbi.2024.07.019. PubMed PMID: 39025418; PubMed Central PMCID: PMC6348822.
 - 25 Sonu S, Post S, Feinglass J. Adverse childhood experiences and the onset of chronic disease in young adulthood. *Prev Med.* 2019;123:163-70. doi: 10.1016/j.ypmed.2019.03.032. PubMed PMID: 30904602.
 - 26 Tan H, Zhou H, Chen J, Ren H, Guo Y, Jiang X. Association of early life adversity with cardiovascular disease and its potential mechanisms: a narrative review. *Front Public Health.* 2024;12:1341266. doi: 10.3389/fpubh.2024.1341266. PubMed PMID: 38362223; PubMed Central PMCID: PMC6348822.

- PMCPMC10867864.
- 27 Friedman EM, Karlamangla AS, Gruenewald TL, Koretz B, Seeman TE. Early life adversity and adult biological risk profiles. *Psychosom Med*. 2015;77:176-85. doi: 10.1097/PSY.000000000000147. PubMed PMID: 25650548; PubMed Central PMCID: PMCPMC4396808.
 - 28 Miller GE, Chen E. Harsh family climate in early life presages the emergence of a proinflammatory phenotype in adolescence. *Psychol Sci*. 2010;21:848-56. doi: 10.1177/0956797610370161. PubMed PMID: 20431047; PubMed Central PMCID: PMCPMC3207635.
 - 29 Miller GE, Chen E, Fok AK, Walker H, Lim A, Nicholls EF, et al. Low early-life social class leaves a biological residue manifested by decreased glucocorticoid and increased proinflammatory signaling. *Proc Natl Acad Sci U S A*. 2009;106:14716-21. doi: 10.1073/pnas.0902971106. PubMed PMID: 19617551; PubMed Central PMCID: PMCPMC2732821.
 - 30 Ehrlich KB, Ross KM, Chen E, Miller GE. Testing the biological embedding hypothesis: Is early life adversity associated with a later proinflammatory phenotype? *Dev Psychopathol*. 2016;28:1273-83. doi: 10.1017/S0954579416000845. PubMed PMID: 27691981; PubMed Central PMCID: PMCPMC5475361.
 - 31 Jung E, Romero R, Yeo L, Diaz-Primera R, Marin-Concha J, Para R, et al. The fetal inflammatory response syndrome: the origins of a concept, pathophysiology, diagnosis, and obstetrical implications. *Semin Fetal Neonatal Med*. 2020;25:101146. doi: 10.1016/j.siny.2020.101146. PubMed PMID: 33164775; PubMed Central PMCID: PMCPMC10580248.
 - 32 Giovannini E, Bonasoni MP, Pascali JP, Giorgetti A, Pelletti G, Gargano G, et al. Infection Induced Fetal Inflammatory Response Syndrome (FIRS): State-of-the-Art and Medico-Legal Implications-A Narrative Review. *Microorganisms*. 2023;11. doi: 10.3390/microorganisms11041010. PubMed PMID: 37110434; PubMed Central PMCID: PMCPMC10142209.
 - 33 Gibson B, Goodfriend E, Zhong Y, Melhem NM. Fetal inflammatory response and risk for psychiatric disorders. *Transl Psychiatry*. 2023;13:224. doi: 10.1038/s41398-023-02505-3. PubMed PMID: 37355708; PubMed Central PMCID: PMCPMC10290670.
 - 34 Lalor MK, Floyd S, Gorak-Stolinska P, Ben-Smith A, Weir RE, Smith SG, et al. BCG vaccination induces different cytokine profiles following infant BCG vaccination in the UK and Malawi. *J Infect Dis*. 2011;204:1075-85. doi: 10.1093/infdis/jir515. PubMed PMID: 21881123; PubMed Central PMCID: PMCPMC3164434.
 - 35 Kagina BM, Abel B, Scriba TJ, Hughes EJ, Keyser A, Soares A, et al. Specific T cell frequency and cytokine expression profile do not correlate with protection against tuberculosis after bacillus Calmette-Guerin vaccination of newborns. *Am J Respir Crit Care Med*. 2010;182:1073-9. doi: 10.1164/rccm.201003-0334OC. PubMed PMID: 20558627; PubMed Central PMCID: PMCPMC2970848.
 - 36 Soares AP, Kwong Chung CK, Choice T, Hughes EJ, Jacobs G, van Rensburg EJ, et al. Longitudinal changes in CD4(+) T-cell memory responses induced by BCG vaccination of newborns. *J Infect Dis*. 2013;207:1084-94. doi: 10.1093/infdis/jis941. PubMed PMID: 23293360; PubMed Central PMCID: PMCPMC3583271.
 - 37 Pablos-Mendez A, Raviglione MC, Laszlo A, Binkin N, Rieder HL, Bustreo F, et al. Global surveillance for antituberculosis-drug resistance, 1994-1997. World Health Organization-International Union against Tuberculosis and Lung Disease Working Group on Anti-Tuberculosis Drug Resistance Surveillance. *N Engl J Med*. 1998;338:1641-9. doi: 10.1056/NEJM199806043382301. PubMed PMID: 9614254.
 - 38 Neurath MF, Finotto S. IL-6 signaling in autoimmunity, chronic inflammation and inflammation-associated cancer. *Cytokine Growth Factor Rev*. 2011;22:83-9. doi: 10.1016/j.cytogfr.2011.02.003. PubMed PMID: 21377916.
 - 39 Wu YQ, Shen J, Zhou QL, Zhao HW, Liu LR, Liu X. Interleukin-6 and interleukin-8 in diagnosing neonatal septicemia. *J Biol Regul Homeost Agents*. 2016;30:1107-13. PubMed PMID: 28078861.
 - 40 Lyra ESNM, Goncalves RA, Pascoal TA, Lima-Filho RAS, Resende EPF, Vieira ELM, et al. Pro-inflammatory interleukin-6 signaling links cognitive impairments and peripheral metabolic alterations in Alzheimer's disease. *Transl Psychiatry*. 2021;11:251. doi: 10.1038/s41398-021-01349-z. PubMed PMID: 33911072; PubMed Central PMCID: PMCPMC8080782.
 - 41 Yao Y, Cai X, Ye Y, Wang F, Chen F, Zheng C. The Role of Microbiota in Infant Health: From Early Life to Adulthood. *Front Immunol*. 2021;12:708472. doi:

- 10.3389/fimmu.2021.708472. PubMed PMID: 34691021; PubMed Central PMCID: PMCPMC8529064.
- 42 Morse ZJ, Simister RL, Crowe SA, Horwitz MS, Osborne LC. Virus induced dysbiosis promotes type 1 diabetes onset. *Front Immunol.* 2023;14:1096323. doi: 10.3389/fimmu.2023.1096323. PubMed PMID: 36742327; PubMed Central PMCID: PMCPMC9892191.
 - 43 Yang L, Jin R, Lu D, Ge Q. T cell Tolerance in Early Life. *Front Immunol.* 2020;11:576261. doi: 10.3389/fimmu.2020.576261. PubMed PMID: 33329542; PubMed Central PMCID: PMCPMC7714828.
 - 44 Ndure J, Flanagan KL. Targeting regulatory T cells to improve vaccine immunogenicity in early life. *Front Microbiol.* 2014;5:477. doi: 10.3389/fmicb.2014.00477. PubMed PMID: 25309517; PubMed Central PMCID: PMCPMC4161046.
 - 45 Batista-Duarte A, Tellez-Martinez D, Fuentes DLP, Carlos IZ. Molecular adjuvants that modulate regulatory T cell function in vaccination: A critical appraisal. *Pharmacol Res.* 2018;129:237-50. doi: 10.1016/j.phrs.2017.11.026. PubMed PMID: 29175113.
 - 46 Bernini L, Manzini CU, Ferri C. BCG and autoimmunity. *Vaccines and Autoimmunity.* 2015:197-206. doi: 10.1002/9781118663721.ch21.
 - 47 Smyk DS, Sakkas LI, Shoenfeld Y, Bogdanos DP. Hepatitis B vaccination and autoimmunity. *Vaccines and Autoimmunity.* 2015:147-62. doi: 10.1002/9781118663721.ch16.
 - 48 Goenka A, Kollmann TR. Development of immunity in early life. *J Infect.* 2015;71:S112-20. doi: 10.1016/j.jinf.2015.04.027. PubMed PMID: 25934325.
 - 49 Huang H, Shu N, Mishra V, Jeon T, Chalak L, Wang ZJ, et al. Development of human brain structural networks through infancy and childhood. *Cereb Cortex.* 2015;25:1389-404. doi: 10.1093/cercor/bht335. PubMed PMID: 24335033; PubMed Central PMCID: PMCPMC4397575.
 - 50 Ercolini AM, Miller SD. The role of infections in autoimmune disease. *Clin Exp Immunol.* 2009;155:1-15. doi: 10.1111/j.1365-2249.2008.03834.x. PubMed PMID: 19076824; PubMed Central PMCID: PMCPMC2665673.
 - 51 Anaya JM, Restrepo-Jimenez P, Ramirez-Santana C. The autoimmune ecology: an update. *Curr Opin Rheumatol.* 2018;30:350-60. doi: 10.1097/BOR.0000000000000498. PubMed PMID: 29438164.
 - 52 Schattner A, Rager-Zisman B. Virus-induced autoimmunity. *Rev Infect Dis.* 1990;12:204-22. doi: 10.1093/clinids/12.2.204. PubMed PMID: 2091619.
 - 53 von Herrath MG, Fujinami RS, Whitton JL. Microorganisms and autoimmunity: making the barren field fertile? *Nat Rev Microbiol.* 2003;1:151-7. doi: 10.1038/nrmicro754. PubMed PMID: 15035044.
 - 54 Cohen AD, Shoenfeld Y. Vaccine-induced autoimmunity. *J Autoimmun.* 1996;9:699-703. doi: 10.1006/jaut.1996.0091. PubMed PMID: 9115571.
 - 55 Agmon-Levin N, Paz Z, Israeli E, Shoenfeld Y. Vaccines and autoimmunity. *Nat Rev Rheumatol.* 2009;5:648-52. doi: 10.1038/nrrheum.2009.196. PubMed PMID: 19865091.
 - 56 Arango M-T, Shoenfeld Y, Cervera R, Anaya J-M. Infection and autoimmune diseases. *Autoimmunity: from bench to bedside.* Bogota: El Rosario University Press; 2013.
 - 57 Geier DA, Geier MR. A case-control study of serious autoimmune adverse events following hepatitis B immunization. *Autoimmunity.* 2005;38:295-301. doi: 10.1080/08916930500144484. PubMed PMID: 16206512.
 - 58 Bogdanos DP, Smith H, Ma Y, Baum H, Mieli-Vergani G, Vergani D. A study of molecular mimicry and immunological cross-reactivity between hepatitis B surface antigen and myelin mimics. *Clin Dev Immunol.* 2005;12:217-24. doi: 10.1080/17402520500285247. PubMed PMID: 16295528; PubMed Central PMCID: PMCPMC2275415.
 - 59 Gherardi RK, Coquet M, Cherin P, Belec L, Moretto P, Dreyfus PA, et al. Macrophagic myofasciitis lesions assess long-term persistence of vaccine-derived aluminium hydroxide in muscle. *Brain.* 2001;124:1821-31. doi: 10.1093/brain/124.9.1821. PubMed PMID: 11522584.
 - 60 Geier MR, Geier DA, Zahalsky AC. A review of hepatitis B vaccination. *Expert Opin Drug Saf.* 2003;2:113-22. doi: 10.1517/14740338.2.2.113. PubMed PMID: 12904111.
 - 61 Maubec E, Pinquier L, Viguier M, Caux F, Amsler E, Aractingi S, et al. Vaccination-induced cutaneous pseudolymphoma. *J Am Acad Dermatol.* 2005;52:623-9. doi: 10.1016/j.jaad.2004.12.021. PubMed PMID: 15793512.
 - 62 Junior DST. Environmental and individual factors associated with protection and predisposition to autoimmune diseases. *Int J Health Sci (Qassim).* 2020;14:13-23. PubMed PMID: 33192227; PubMed Central

- PMCID: PMCPMC7644454.
- 63 Ortega-Hernandez OD, Shoenfeld Y. Infection, vaccination, and autoantibodies in chronic fatigue syndrome, cause or coincidence? *Ann N Y Acad Sci.* 2009;1173:600-9. doi: 10.1111/j.1749-6632.2009.04799.x. PubMed PMID: 19758205.
 - 64 Mikaeloff Y, Caridade G, Suissa S, Tardieu M. Hepatitis B vaccine and the risk of CNS inflammatory demyelination in childhood. *Neurology.* 2009;72:873-80. doi: 10.1212/01.wnl.0000335762.42177.07. PubMed PMID: 18843097.
 - 65 Herroelen L, de Keyser J, Ebinger G. Central-nervous-system demyelination after immunisation with recombinant hepatitis B vaccine. *Lancet.* 1991;338:1174-5. doi: 10.1016/0140-6736(91)92034-y. PubMed PMID: 1682594.
 - 66 Tourbah A, Gout O, Liblau R, Lyon-Caen O, Boungniot C, Iba-Zizen MT, et al. Encephalitis after hepatitis B vaccination: recurrent disseminated encephalitis or MS? *Neurology.* 1999;53:396-401. doi: 10.1212/wnl.53.2.396. PubMed PMID: 10430433.
 - 67 Agmon-Levin N, Zafrir Y, Kivity S, Balofsky A, Amital H, Shoenfeld Y. Chronic fatigue syndrome and fibromyalgia following immunization with the hepatitis B vaccine: another angle of the 'autoimmune (auto-inflammatory) syndrome induced by adjuvants' (ASIA). *Immunol Res.* 2014;60:376-83. doi: 10.1007/s12026-014-8604-2. PubMed PMID: 25427994.
 - 68 Nancy AL, Shoenfeld Y. Chronic fatigue syndrome with autoantibodies--the result of an augmented adjuvant effect of hepatitis-B vaccine and silicone implant. *Autoimmun Rev.* 2008;8:52-5. doi: 10.1016/j.autrev.2008.07.026. PubMed PMID: 18725327.
 - 69 Richardson CT, Hayden MS, Gilmore ES, Poligone B. Evaluation of the Relationship between Alopecia Areata and Viral Antigen Exposure. *Am J Clin Dermatol.* 2018;19:119-26. doi: 10.1007/s40257-017-0312-y. PubMed PMID: 28801732.
 - 70 Choffray A, Piquier L, Bachelez H. Exacerbation of lupus panniculitis following anti-hepatitis-B vaccination. *Dermatology.* 2007;215:152-4. doi: 10.1159/000104268. PubMed PMID: 17684379.
 - 71 Luhadia K, Abugrin M, Kiani R, Ahmady A, Virk J, Yashi K. Hepatitis B Vaccine and Multiple Sclerosis: Cause or Coincidence. *Cureus.* 2022;14:e29941. doi: 10.7759/cureus.29941. PubMed PMID: 36348919; PubMed Central PMCID: PMCPMC9635015.
 - 72 Limas C, Limas CJ. Lichen planus in children: a possible complication of hepatitis B vaccines. *Pediatr Dermatol.* 2002;19:204-9. doi: 10.1046/j.1525-1470.2002.00079.x. PubMed PMID: 12047638.
 - 73 de la Fuente S, Hernandez-Martin A, de Lucas R, Gonzalez-Ensenat MA, Vicente A, Colmenero I, et al. Postvaccination bullous pemphigoid in infancy: report of three new cases and literature review. *Pediatr Dermatol.* 2013;30:741-4. doi: 10.1111/pde.12231. PubMed PMID: 24125034.
 - 74 Erbagci Z. Childhood bullous pemphigoid following hepatitis B immunization. *J Dermatol.* 2002;29:781-5. doi: 10.1111/j.1346-8138.2002.tb00223.x. PubMed PMID: 12532044.
 - 75 Berkun Y, Mimouni D, Shoenfeld Y. Pemphigus following hepatitis B vaccination--coincidence or causality? *Autoimmunity.* 2005;38:117-9. doi: 10.1080/08916930400027078. PubMed PMID: 16040331.
 - 76 Vital C, Vital A, Gbikpi-Benissan G, Longy-Boursier M, Climas MT, Castaing Y, et al. Postvaccinal inflammatory neuropathy: peripheral nerve biopsy in 3 cases. *J Peripher Nerv Syst.* 2002;7:163-7. doi: 10.1046/j.1529-8027.2002.02010.x. PubMed PMID: 12365564.
 - 77 de Carvalho JF, Pereira RM, Shoenfeld Y. Systemic polyarteritis nodosa following hepatitis B vaccination. *Eur J Intern Med.* 2008;19:575-8. doi: 10.1016/j.ejim.2007.06.035. PubMed PMID: 19046721.
 - 78 Mailliefert JF, Farge P, Gazet-Maillefert MP, Tavernier C. Mental nerve neuropathy as a result of hepatitis B vaccination. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 1997;83:663-4. doi: 10.1016/s1079-2104(97)90315-2. PubMed PMID: 9195619.
 - 79 Zaas A, Scheel P, Venbrux A, Hellmann DB. Large artery vasculitis following recombinant hepatitis B vaccination: 2 cases. *J Rheumatol.* 2001;28:1116-20. PubMed PMID: 11361200.
 - 80 Agmon-Levin N, Zafrir Y, Paz Z, Shilton T, Zandman-Goddard G, Shoenfeld Y. Ten cases of systemic lupus erythematosus related to hepatitis B vaccine. *Lupus.* 2009;18:1192-7. doi: 10.1177/0961203309345732. PubMed PMID: 19880567.
 - 81 Altman A, Szyper-Kravitz M, Shoenfeld Y. HBV vaccine and dermatomyositis: is there an association? *Rheumatol Int.* 2008;28:609-12. doi: 10.1007/s00296-007-0485-4. PubMed PMID: 18034245.
 - 82 Geier MR, Geier DA. A case-series of

- adverse events, positive re-challenge of symptoms, and events in identical twins following hepatitis B vaccination: analysis of the Vaccine Adverse Event Reporting System (VAERS) database and literature review. *Clin Exp Rheumatol*. 2004;22:749-55. PubMed PMID: 15638050.
- 83 Geier MR, Geier DA. Hepatitis B vaccination safety. *Ann Pharmacother*. 2002;36:370-4. doi: 10.1345/aph.1A304. PubMed PMID: 11895045.
 - 84 Pennesi M, Torre G, Del Santo M, Sonzogni A. Glomerulonephritis after recombinant hepatitis B vaccine. *Pediatr Infect Dis J*. 2002;21:172-3. doi: 10.1097/00006454-200202000-00020. PubMed PMID: 11840090.
 - 85 Poirriez J. A preliminary experiment of absorption of antinuclear antibodies by the hepatitis B vaccine components, in a case of neurolupus. *Vaccine*. 2004;22:3166-8. doi: 10.1016/j.vaccine.2004.01.056. PubMed PMID: 15297069.
 - 86 Schattner A. Consequence or coincidence? The occurrence, pathogenesis and significance of autoimmune manifestations after viral vaccines. *Vaccine*. 2005;23:3876-86. doi: 10.1016/j.vaccine.2005.03.005. PubMed PMID: 15917108.
 - 87 Ramirez-Rivera J, Vega-Cruz AM, Jaime-Anselmi F. Polymyositis: rare complication of hepatitis B vaccination. An unusual cause of toxic shock syndrome. *Bol Asoc Med P R*. 2003;95:13-6. PubMed PMID: 15449786.
 - 88 Agmon-Levin N, Kivity S, Szyper-Kravitz M, Shoenfeld Y. Transverse myelitis and vaccines: a multi-analysis. *Lupus*. 2009;18:1198-204. doi: 10.1177/0961203309345730. PubMed PMID: 19880568.
 - 89 Maillefert JF, Sibilia J, Toussiroit E, Vignon E, Eschard JP, Lorcier B, et al. Rheumatic disorders developed after hepatitis B vaccination. *Rheumatology (Oxford)*. 1999;38:978-83. doi: 10.1093/rheumatology/38.10.978. PubMed PMID: 10534549.
 - 90 Ronchi F, Cecchi P, Falcioni F, Marsciani A, Minak G, Muratori G, et al. Thrombocytopenic purpura as adverse reaction to recombinant hepatitis B vaccine. *Arch Dis Child*. 1998;78:273-4. doi: 10.1136/adc.78.3.273. PubMed PMID: 9613364; PubMed Central PMCID: PMC1717498.
 - 91 Neau D, Bonnet F, Michaud M, Perel Y, Longy-Boursier M, Ragnaud JM, et al. Immune thrombocytopenic purpura after recombinant hepatitis B vaccine: retrospective study of seven cases. *Scand J Infect Dis*. 1998;30:115-8. doi: 10.1080/003655498750003465. PubMed PMID: 9730294.
 - 92 Chave T, Neal C, Camp R. Henoch-Schönlein purpura following hepatitis B vaccination. *J Dermatolog Treat*. 2003;14:179-81. doi: 10.1080/09546630310004199. PubMed PMID: 14522629.
 - 93 Khamaisi M, Shoenfeld Y, Orbach H, Guillain-Barre syndrome following hepatitis B vaccination. *Clin Exp Rheumatol*. 2004;22:767-70. PubMed PMID: 15638054.
 - 94 Girard M. Autoimmune hazards of hepatitis B vaccine. *Autoimmun Rev*. 2005;4:96-100. doi: 10.1016/j.autrev.2004.10.002. PubMed PMID: 15722255.
 - 95 Wise RP, Kiminyo KP, Salive ME. Hair loss after routine immunizations. *JAMA*. 1997;278:1176-8. PubMed PMID: 9326478.
 - 96 Avci Z, Bayram C, Malbora B. Hepatitis B vaccine-associated atypical hemolytic uremic syndrome. *Turk J Haematol*. 2013;30:418-9. doi: 10.4274/Tjh-2013.0226. PubMed PMID: 24385836; PubMed Central PMCID: PMC3874978.
 - 97 Edelman R. The development and use of vaccine adjuvants. *Mol Biotechnol*. 2002;21:129-48. doi: 10.1385/MB:21:2:129. PubMed PMID: 12059113.
 - 98 Pulendran B, Ahmed R. Immunological mechanisms of vaccination. *Nat Immunol*. 2011;12:509-17. doi: 10.1038/ni.2039. PubMed PMID: 21739679; PubMed Central PMCID: PMC3253344.
 - 99 Kuroda E, Coban C, Ishii KJ. Particulate adjuvant and innate immunity: past achievements, present findings, and future prospects. *Int Rev Immunol*. 2013;32:209-20. doi: 10.3109/08830185.2013.773326. PubMed PMID: 23570316; PubMed Central PMCID: PMC3632271.
 - 100 Guimaraes LE, Baker B, Perricone C, Shoenfeld Y. Vaccines, adjuvants and autoimmunity. *Pharmacol Res*. 2015;100:190-209. doi: 10.1016/j.phrs.2015.08.003. PubMed PMID: 26275795; PubMed Central PMCID: PMC3632276.
 - 101 Boretti A. Reviewing the association between aluminum adjuvants in the vaccines and autism spectrum disorder. *J Trace Elem Med Biol*. 2021;66:126764. doi: 10.1016/j.jtemb.2021.126764. PubMed PMID: 33930617.
 - 102 Shoenfeld Y, Agmon-Levin N. 'ASIA' - autoimmune/inflammatory syndrome induced by adjuvants. *J Autoimmun*. 2011;36:4-8. doi: 10.1016/j.jaut.2010.07.003. PubMed PMID: 20708902.
 - 103 Zafrir Y, Agmon-Levin N, Paz Z, Shilton T, Shoenfeld Y. Autoimmunity following

- hepatitis B vaccine as part of the spectrum of 'Autoimmune (Auto-inflammatory) Syndrome induced by Adjuvants' (ASIA): analysis of 93 cases. *Lupus*. 2012;21:146-52. doi: 10.1177/0961203311429318. PubMed PMID: 22235045.
- 104 Shaw CA, Tomljenovic L. Aluminum in the central nervous system (CNS): toxicity in humans and animals, vaccine adjuvants, and autoimmunity. *Immunol Res*. 2013;56:304-16. doi: 10.1007/s12026-013-8403-1. PubMed PMID: 23609067.
- 105 Ewing GE. What is regressive autism and why does it occur? Is it the consequence of multi-systemic dysfunction affecting the elimination of heavy metals and the ability to regulate neural temperature? *N Am J Med Sci*. 2009;1:28-47. PubMed PMID: 22666668; PubMed Central PMCID: PMCPMC3364648.
- 106 Tomljenovic L, Shaw CA. Aluminum vaccine adjuvants: are they safe? *Curr Med Chem*. 2011;18:2630-7. doi: 10.2174/092986711795933740. PubMed PMID: 21568886.
- 107 Borba V, Malkova A, Basantsova N, Halpert G, Andreoli L, Tincani A, et al. Classical Examples of the Concept of the ASIA Syndrome. *Biomolecules*. 2020;10. doi: 10.3390/biom10101436. PubMed PMID: 33053910; PubMed Central PMCID: PMCPMC7600067.
- 108 Marciano BE, Huang CY, Joshi G, Rezaei N, Carvalho BC, Allwood Z, et al. BCG vaccination in patients with severe combined immunodeficiency: complications, risks, and vaccination policies. *J Allergy Clin Immunol*. 2014;133:1134-41. doi: 10.1016/j.jaci.2014.02.028. PubMed PMID: 24679470; PubMed Central PMCID: PMCPMC4015464.
- 109 Jain M, Vadboncoeur J, Garg SJ, Biswas J. Bacille Calmette-Guerin: An ophthalmic perspective. *Surv Ophthalmol*. 2022;67:307-20. doi: 10.1016/j.survophthal.2021.07.005. PubMed PMID: 34343536; PubMed Central PMCID: PMCPMC8325561.
- 110 Khalili N, Mohammadzadeh I, Khalili N, Heredia RJ, Zoghi S, Boztug K, et al. BCGitis as the primary manifestation of chronic granulomatous disease. *IDCases*. 2021;23:e01038. doi: 10.1016/j.idcr.2020.e01038. PubMed PMID: 33425681; PubMed Central PMCID: PMCPMC7785948.
- 111 Dahl J, Holvik K, Heldal E, Grimnes G, Hoff M, Finnes TE, et al. Individual Variation in Adaptive Immune Responses and Risk of Hip Fracture-A NOREPOS Population-Based Cohort Study. *J Bone Miner Res*. 2020;35:2327-34. doi: 10.1002/jbmr.4135. PubMed PMID: 32697001.
- 112 Tsujioka Y, Nozaki T, Nishimura G, Miyazaki O, Jinzaki M, Kono T. BCG osteomyelitis: tips for diagnosis. *Skeletal Radiol*. 2022;51:1571-84. doi: 10.1007/s00256-021-03966-7. PubMed PMID: 35043224.
- 113 Wang J, Zhou F, Jiang MB, Xu ZH, Ni YH, Wu QS. Epidemiological characteristics and trends of Bacillus Calmette-Guerin lymphadenitis in Shanghai, China from 2010 to 2019. *Hum Vaccin Immunother*. 2022;18:1938922. doi: 10.1080/21645515.2021.1938922. PubMed PMID: 34156902; PubMed Central PMCID: PMCPMC8920159.
- 114 Sellami K, Amouri M, Kmiha S, Bahloul E, Aloulou H, Sfaihi L, et al. Adverse Reactions Due to the Bacillus Calmette-Guerin Vaccine: Twenty Tunisian Cases. *Indian J Dermatol*. 2018;63:62-5. doi: 10.4103/ij.d.IJD_327_17. PubMed PMID: 29527028; PubMed Central PMCID: PMCPMC5838757.
- 115 Modrzejewska M, Karczewicz D, Kordek A, Rudnicki J, Czajka R. [The detachment of retina as possible complication after BCG vaccination during HOP--description of case]. *Klin Oczna*. 2006;108:446-9. PubMed PMID: 17455724.
- 116 Salmon C, Conus F, Parent ME, Benedetti A, Rousseau MC. Association between Bacillus Calmette-Guerin vaccination and lymphoma: a population-based birth cohort study. *J Intern Med*. 2019;286:583-95. doi: 10.1111/joim.12965. PubMed PMID: 31361936.
- 117 Shoenfeld Y, Aron-Maor A, Tanai A, Ehrenfeld M. Bcg and autoimmunity: another two-edged sword. *J Autoimmun*. 2001;16:235-40. doi: 10.1006/jaut.2000.0494. PubMed PMID: 11334488.
- 118 Schuchmann L, Pernice W, Hufschmidt C, Adler CP. Tuberculous arthritis--a rare, but important differential diagnosis in juvenile chronic arthritis. *Monatsschr Kinderheilkd*. 1991;139:244-7. PubMed PMID: 2072968.
- 119 Anis O, Yogev D, Dotan A, Tsur AM, David P, Vishnevskia Dai V, et al. Autoimmune disorders caused by intravesical bacillus Calmette-Guerine treatment: A systematic review. *Autoimmun Rev*. 2023;22:103329. doi: 10.1016/j.autrev.2023.103329. PubMed PMID: 37061015.
- 120 Sharan S, Thirkill CE, Grigg JR. Autoimmune retinopathy associated with intravesical BCG therapy. *Br J Ophthalmol*. 2005;89:927-8. doi: 10.1136/bjo.2004.065359. PubMed PMID: 15965187; PubMed Central PMCID: PMCPMC1772734.
- 121 Nakagawa T, Shigehara K, Naito R, Yaegashi H, Nakashima K, Iijima M, et al. Reiter's syndrome following intravesical Bacillus

- Calmette-Guerin therapy for bladder carcinoma: a report of five cases. *Int Cancer Conf J*. 2018;7:148-51. doi: 10.1007/s13691-018-0342-1. PubMed PMID: 31149535; PubMed Central PMCID: PMC6498336.
- 122 Genereau T, Koeger AC, Chaibi P, Bourgeois P. Polymyalgia rheumatica with temporal arteritis following intravesical Calmette-Guerin bacillus immunotherapy for bladder cancer. *Clin Exp Rheumatol*. 1996;14:110. PubMed PMID: 8697650.
- 123 Thepot C, Martigny J, Simon L, Bellin J, Larget-Piet B, Chevalier X. Acute polyarthritis after BCG-therapy for bladder carcinoma in a patient with ankylosing spondylitis. *Rev Rhum Engl Ed*. 1995;62:459-61. PubMed PMID: 7552212.
- 124 Granel B, Serratrice J, Morange PE, Disdier P, Weiller PJ. Cryoglobulinemia vasculitis following intravesical instillations of bacillus Calmette-Guerin. *Clin Exp Rheumatol*. 2004;22:481-2. PubMed PMID: 15301249.
- 125 Tsuchiya H, Hanata N, Harada H, Shoda H, Fujio K. Intestinal ulcers induced by intravesical bacillus Calmette-Guerin therapy. *Mod Rheumatol Case Rep*. 2021;5:421-4. doi: 10.1080/24725625.2021.1877381. PubMed PMID: 33492192.
- 126 Parent ME, Richer M, Liang P. The first case of bacillus Calmette-Guerin-induced small-vessel central nervous system vasculitis. *Clin Rheumatol*. 2018;37:2297-302. doi: 10.1007/s10067-018-4136-9. PubMed PMID: 29740727.
- 127 Beisland C, Holsen DS. Vitiligo--an autoimmune side-effect of intravesical bacillus Calmette-Guerin instillation? *Scand J Urol Nephrol*. 2004;38:182-3. doi: 10.1080/00365590310021357. PubMed PMID: 15204415.
- 128 Vittori F, Gros-lafeige C. [Tuberculosis lupus after BCG vaccination. A rare complication of the vaccination]. *Arch Pediatr*. 1996;3:457-9. doi: 10.1016/0929-693x(96)86404-9. PubMed PMID: 8763716.
- 129 Izumi AK, Matsunaga J. BCG vaccine-induced lupus vulgaris. *Arch Dermatol*. 1982;118:171-2. PubMed PMID: 7065666.
- 130 Noishiki C, Hayasaka Y, Yoshida R, Ogawa R. Over 90% of Childhood BCG Vaccine-Induced Keloids in Japan Occur in Women. *Dermatol Ther (Heidelb)*. 2023;13:1137-47. doi: 10.1007/s13555-023-00916-0. PubMed PMID: 36952124; PubMed Central PMCID: PMC6410034230.
- 131 Abid H, Fiquigui M, Adil Ibrahim S, El Abkari M, Mzyiene M, Ennaciri S, et al. Acute Hepatitis Induced by Intravesical BCG Therapy: A Rare but Serious Complication. *Case Reports Hepatol*. 2021;2021:4574879. doi: 10.1155/2021/4574879. PubMed PMID: 34258085; PubMed Central PMCID: PMC6410034230.
- 132 Yamazaki-Nakashimada MA, Unzueta A, Berenise Gamez-Gonzalez L, Gonzalez-Saldana N, Sorensen RU. BCG: a vaccine with multiple faces. *Hum Vaccin Immunother*. 2020;16:1841-50. doi: 10.1080/21645515.2019.1706930. PubMed PMID: 31995448; PubMed Central PMCID: PMC6410034230.
- 133 Sumida Y, Kanemasa K, Tachibana S, Maekawa K, Nakano T. A case of autoimmune pancreatitis occurring during intravesical bacillus Calmette Guerin immunotherapy for ureteral cancer. *Nihon Shokakibyō Gakkai Zasshi*. 2003;100:1328-32. PubMed PMID: 14639932.
- 134 Foucard T, Hjelmstedt A. BCG-osteomyelitis and -osteoarthritis as a complication following BCG-vaccination. *Acta Orthop Scand*. 1971;42:142-51. doi: 10.3109/17453677108989034. PubMed PMID: 4939582.
- 135 Zawar V, Chuh A. A case-control study on the association of pulse oral poliomyelitis vaccination and Gianotti-Crosti syndrome. *Int J Dermatol*. 2017;56:75-9. doi: 10.1111/ijd.13451. PubMed PMID: 27943306.
- 136 Gao J, Kang G, Hu R, Zhang L, Yu J, Wang Z, et al. Adverse events following immunization with bivalent oral poliovirus vaccine in Jiangsu, China. *Br J Clin Pharmacol*. 2021;87:4831-8. doi: 10.1111/bcp.14976. PubMed PMID: 34240463.
- 137 Akbayram S, Karaman K, Ece I, Hatice Akbayram T. Acute immune thrombocytopenic purpura following oral polio vaccination. *Platelets*. 2015;26:705. doi: 10.3109/09537104.2014.980798. PubMed PMID: 25384121.
- 138 Elkhayat HA, El-Rashidy OF, Elagouza IA, Zaitoun R, Abbas YAA. Childhood acute disseminated encephalomyelitis: an Egyptian pilot study. *Acta Neurol Belg*. 2020;120:549-55. doi: 10.1007/s13760-017-0869-5. PubMed PMID: 29222716.
- 139 Kelly H. Evidence for a causal association between oral polio vaccine and transverse myelitis: A case history and review of the Literature. *J Paediatr Child Health*. 2006;42:155-9. doi: 10.1111/j.1440-1754.2006.00840.x. PubMed PMID: 16630313.
- 140 Hughes AM, Ponsonby AL, Dear K, Dwyer T, Taylor BV, van der Mei I, et al. Childhood infections, vaccinations, and tonsillectomy

- and risk of first clinical diagnosis of CNS demyelination in the Ausimmune Study. *Mult Scler Relat Disord.* 2020;42:102062. doi: 10.1016/j.msard.2020.102062. PubMed PMID: 32305688.
- 141 Pineton de Chambrun G, Dauchet L, Gower-Rousseau C, Cortot A, Colombel JF, Peyrin-Biroulet L. Vaccination and Risk for Developing Inflammatory Bowel Disease: A Meta-Analysis of Case-Control and Cohort Studies. *Clin Gastroenterol Hepatol.* 2015;13:1405-15. doi: 10.1016/j.cgh.2015.04.179. PubMed PMID: 25956840.
- 142 Platt LR, Estivariz CF, Sutter RW. Vaccine-associated paralytic poliomyelitis: a review of the epidemiology and estimation of the global burden. *J Infect Dis.* 2014;210:S380-9. doi: 10.1093/infdis/jiu184. PubMed PMID: 25316859; PubMed Central PMCID: PMC424844.
- 143 Dhiman R, Prakash SC, Sreenivas V, Puliyl J. Correlation between Non-Polio Acute Flaccid Paralysis Rates with Pulse Polio Frequency in India. *Int J Environ Res Public Health.* 2018;15. doi: 10.3390/ijerph15081755. PubMed PMID: 30111741; PubMed Central PMCID: PMC6121585.
- 144 Puliyl J, Naik P. Revised World Health Organization (WHO)'s causality assessment of adverse events following immunization-a critique. *F1000Res.* 2018;7:243. doi: 10.12688/f1000research.13694.2. PubMed PMID: 30026925; PubMed Central PMCID: PMC6039921.
- 145 Hooker BS, Miller NZ. Analysis of health outcomes in vaccinated and unvaccinated children: Developmental delays, asthma, ear infections and gastrointestinal disorders. *SAGE Open Med.* 2020;8:2050312120925344. doi: 10.1177/2050312120925344. PubMed PMID: 32537156; PubMed Central PMCID: PMC7268563.
- 146 Hooker BS, Miller NZ. Health effects in vaccinated versus unvaccinated children, with covariates for breastfeeding status and type of birth. *Journal of Translational Science.* 2021;7:1-11. doi: 10.15761/JTS.1000459.
- 147 Mawson AR, Ray BD, Bhuiyan AR, Jacob B. Pilot comparative study on the health of vaccinated and unvaccinated 6-to 12-year-old US children. *J Transl Sci.* 2017;3:1-12. doi: 10.15761/JTS.1000186.
- 148 Basha S, Surendran N, Pichichero M. Immune responses in neonates. *Expert Rev Clin Immunol.* 2014;10:1171-84. doi: 10.1586/1744666X.2014.942288. PubMed PMID: 25088080; PubMed Central PMCID: PMC4407563.
- 149 Marchant A, Kollmann TR. Understanding the ontogeny of the immune system to promote immune-mediated health for life. *Front Immunol.* 2015;6:77. doi: 10.3389/fimmu.2015.00077. PubMed PMID: 25755655; PubMed Central PMCID: PMC437332.
- 150 Adkins B, Leclerc C, Marshall-Clarke S. Neonatal adaptive immunity comes of age. *Nat Rev Immunol.* 2004;4:553-64. doi: 10.1038/nri1394. PubMed PMID: 15229474.
- 151 Willems F, Vollstedt S, Suter M. Phenotype and function of neonatal DC. *Eur J Immunol.* 2009;39:26-35. doi: 10.1002/eji.200838391. PubMed PMID: 19137537.
- 152 Leiber A, Graf B, Spring B, Rudner J, Kostlin N, Orlikowsky TW, et al. Neonatal monocytes express antiapoptotic pattern of Bcl-2 proteins and show diminished apoptosis upon infection with *Escherichia coli*. *Pediatr Res.* 2014;76:142-9. doi: 10.1038/pr.2014.74. PubMed PMID: 24850312.
- 153 Abughali N, Berger M, Tosi MF. Deficient total cell content of CR3 (CD11b) in neonatal neutrophils. *Blood.* 1994;83:1086-92. PubMed PMID: 7906565.
- 154 Guilmo A, Hermann E, Braud VM, Carlier Y, Truysens C. Natural killer cell responses to infections in early life. *J Innate Immun.* 2011;3:280-8. doi: 10.1159/000323934. PubMed PMID: 21411972.
- 155 Le Garff-Tavernier M, Beziat V, Decocq J, Siguret V, Gandjbakhch F, Pautas E, et al. Human NK cells display major phenotypic and functional changes over the life span. *Aging Cell.* 2010;9:527-35. doi: 10.1111/j.1474-9726.2010.00584.x. PubMed PMID: 20477761.
- 156 Levy O. Innate immunity of the newborn: basic mechanisms and clinical correlates. *Nat Rev Immunol.* 2007;7:379-90. doi: 10.1038/nri2075. PubMed PMID: 17457344.
- 157 Soulie JC, Devillier P, Santarelli J, Goudeau A, Vermeulen P, Guellier M, et al. Immunogenicity and safety in newborns of a new recombinant hepatitis B vaccine containing the S and pre-S2 antigens. *Vaccine.* 1991;9:545-8. doi: 10.1016/0264-410x(91)90240-7. PubMed PMID: 1837646.
- 158 Strandmark J, Darboe A, Diray-Arce J, Ben-Othman R, Vignolo SM, Rao S, et al. A single birth dose of Hepatitis B vaccine induces polyfunctional CD4(+) T helper cells. *Front Immunol.* 2022;13:1043375. doi: 10.3389/fimmu.2022.1043375. PubMed PMID: 36426360; PubMed Central PMCID: PMC9681035.
- 159 Martinez L, Cords O, Liu Q, Acuna-Villaorduna

- C, Bonnet M, Fox GJ, et al. Infant BCG vaccination and risk of pulmonary and extra-pulmonary tuberculosis throughout the life course: a systematic review and individual participant data meta-analysis. *Lancet Glob Health*. 2022;10:e1307-e16. doi: 10.1016/S2214-109X(22)00283-2. PubMed PMID: 35961354; PubMed Central PMCID: PMCPMC10406427.
- 160 Roy A, Eisenhut M, Harris RJ, Rodrigues LC, Sridhar S, Habermann S, et al. Effect of BCG vaccination against Mycobacterium tuberculosis infection in children: systematic review and meta-analysis. *BMJ*. 2014;349:g4643. doi: 10.1136/bmj.g4643. PubMed PMID: 25097193; PubMed Central PMCID: PMCPMC4122754
- 161 Waggle Z, Geldenhuys H, Sutter RW, Jacks M, Mulenga H, Mahomed H, et al. Randomized trial of type 1 and type 3 oral monovalent poliovirus vaccines in newborns in Africa. *J Infect Dis*. 2012;205:228-36. doi: 10.1093/infdis/jir721. PubMed PMID: 22158680.
- 162 Sutter RW, John TJ, Jain H, Agarkhedkar S, Ramanan PV, Verma H, et al. Immunogenicity of bivalent types 1 and 3 oral poliovirus vaccine: a randomised, double-blind, controlled trial. *Lancet*. 2010;376:1682-8. doi: 10.1016/S0140-6736(10)61230-5. PubMed PMID: 20980048.
- 163 el-Sayed N, el-Gamal Y, Abbassy AA, Seoud I, Salama M, Kandeel A, et al. Monovalent type 1 oral poliovirus vaccine in newborns. *N Engl J Med*. 2008;359:1655-65. doi: 10.1056/NEJMoa0800390. PubMed PMID: 18923170.
- 164 Bhaskaram P, Nair KM, Hemalatha P, Murthy N, Nair P. Systemic and mucosal immune response to polio vaccination with additional dose in newborn period. *J Trop Pediatr*. 1997;43:232-4. doi: 10.1093/tropej/43.4.232. PubMed PMID: 9283127.
- 165 Jain PK, Dutta AK, Nangia S, Khare S, Saili A. Seroconversion following killed polio vaccine in neonates. *Indian J Pediatr*. 1997;64:511-5. doi: 10.1007/BF02737758. PubMed PMID: 10771880.
- 166 Khare S, Kumari S, Nagpal IS, Sharma D, Verghese T. Oral polio vaccination in infants: beneficial effect of additional dose at birth. *Indian J Pediatr*. 1993;60:275-81. doi: 10.1007/BF02822191. PubMed PMID: 8244504.
- 167 Dong DX, Hu XM, Liu WJ, Li JS, Jin YC, Tan SG, et al. Immunization of neonates with trivalent oral poliomyelitis vaccine (Sabin). *Bull World Health Organ*. 1986;64:853-60. PubMed PMID: 3493854; PubMed Central PMCID: PMCPMC2490980.
- 168 Szmunes W, Stevens CE, Harley EJ, Zang EA, Oleszko WR, William DC, et al. Hepatitis B vaccine: demonstration of efficacy in a controlled clinical trial in a high-risk population in the United States. *N Engl J Med*. 1980;303:833-41. doi: 10.1056/NEJM198010093031501. PubMed PMID: 6997738.
- 169 Szmunes W, Stevens CE, Zang EA, Harley EJ, Kellner A. A controlled clinical trial of the efficacy of the hepatitis B vaccine (Heptavax B): a final report. *Hepatology*. 1981;1:377-85. doi: 10.1002/hep.1840010502. PubMed PMID: 7030902.
- 170 Francis DP, Hadler SC, Thompson SE, Maynard JE, Ostrow DG, Altman N, et al. The prevention of hepatitis B with vaccine. Report of the centers for disease control multi-center efficacy trial among homosexual men. *Ann Intern Med*. 1982;97:362-6. doi: 10.7326/0003-4819-97-3-362. PubMed PMID: 6810736.
- 171 Immunisation against hepatitis B. *Lancet*. 1988;1:875-6. PubMed PMID: 2895375.
- 172 Venters C, Graham W, Cassidy W. Recombivax-HB: perspectives past, present and future. *Expert Rev Vaccines*. 2004;3:119-29. doi: 10.1586/14760584.3.2.119. PubMed PMID: 15056038.
- 173 Assad S, Francis A. Over a decade of experience with a yeast recombinant hepatitis B vaccine. *Vaccine*. 1999;18:57-67. doi: 10.1016/s0264-410x(99)00179-6. PubMed PMID: 10501235.
- 174 Stevens CE, Toy PT, Taylor PE, Lee T, Yip HY. Prospects for control of hepatitis B virus infection: implications of childhood vaccination and long-term protection. *Pediatrics*. 1992;90:170-3. PubMed PMID: 1534885.
- 175 Gelinas L, Abu-Raya B, Ruck C, Cai B, Kollmann TR. Hepatitis B virus vaccine-induced cell-mediated immunity correlates with humoral immune response following primary vaccination during infancy. *ImmunoHorizons*. 2017;1:42-52. doi: 10.4049/immunohorizons.1700015.
- 176 Larsen CE, Xu J, Lee S, Dubey DP, Uko G, Yunis EJ, et al. Complex cytokine responses to hepatitis B surface antigen and tetanus toxoid in responders, nonresponders and subjects naive to hepatitis B surface antigen. *Vaccine*. 2000;18:3021-30. doi: 10.1016/s0264-410x(00)00084-0. PubMed PMID: 10825606.
- 177 Jarrosson L, Kolopp-Sarda MN, Aguilar P, Bene MC, Lepori ML, Vignaud MC, et al. Most humoral non-responders to hepatitis

- B vaccines develop HBV-specific cellular immune responses. *Vaccine*. 2004;22:3789-96. doi: 10.1016/j.vaccine.2004.02.046. PubMed PMID: 15315860.
- 178 del Canho R, Grosheide PM, Voogd M, Huisman WM, Heijtkink RA, Schalm SW. Immunogenicity of 20 micrograms of recombinant DNA hepatitis B vaccine in healthy neonates: a comparison of three different vaccination schemes. *J Med Virol*. 1993;41:30-4. doi: 10.1002/jmv.1890410107. PubMed PMID: 8228934.
- 179 Mazel JA, Schalm SW, de Gast BC, Nuijten AS, Heijtkink RA, Botman MJ, et al. Passive-active immunisation of neonates of HBsAg positive carrier mothers: preliminary observations. *Br Med J (Clin Res Ed)*. 1984;288:513-5. doi: 10.1136/bmj.288.6416.513. PubMed PMID: 6421359; PubMed Central PMCID: PMCPMC1444606.
- 180 Schalm SW, Mazel JA, de Gast GC, Heijtkink RA, Botman MJ, Banffer JR, et al. Prevention of hepatitis B infection in newborns through mass screening and delayed vaccination of all infants of mothers with hepatitis B surface antigen. *Pediatrics*. 1989;83:1041-8. PubMed PMID: 2726331.
- 181 Marchant A, Newport M. Prevention of infectious diseases by neonatal and early infantile immunization: prospects for the new millennium. *Curr Opin Infect Dis*. 2000;13:241-6. doi: 10.1097/00001432-200006000-00007. PubMed PMID: 11964793.
- 182 Siegrist CA. Neonatal and early life vaccinology. *Vaccine*. 2001;19:3331-46. doi: 10.1016/s0264-410x(01)00028-7. PubMed PMID: 11348697.
- 183 Marchant A, Pihlgren M, Goetghebuer T, Weiss HA, Ota MO, Schlegel-Hauter SE, et al. Predominant influence of environmental determinants on the persistence and avidity maturation of antibody responses to vaccines in infants. *J Infect Dis*. 2006;193:1598-605. doi: 10.1086/503775. PubMed PMID: 16652290.
- 184 Ansari A, Vincent JP, Moorhouse L, Shimakawa Y, Nayagam S. Risk of early horizontal transmission of hepatitis B virus in children of uninfected mothers in sub-Saharan Africa: a systematic review and meta-analysis. *Lancet Glob Health*. 2023;11:e715-e28. doi: 10.1016/S2214-109X(23)00131-6. PubMed PMID: 37061310.
- 185 Tena-Coki NG, Scriba TJ, Peteni N, Eley B, Wilkinson RJ, Andersen P, et al. CD4 and CD8 T-cell responses to mycobacterial antigens in African children. *Am J Respir Crit Care Med*. 2010;182:120-9. doi: 10.1164/rccm.200912-1862OC. PubMed PMID: 20224065; PubMed Central PMCID: PMCPMC2902756.
- 186 Kagina BM, Abel B, Bowmaker M, Scriba TJ, Gelderbloem S, Smit E, et al. Delaying BCG vaccination from birth to 10 weeks of age may result in an enhanced memory CD4 T cell response. *Vaccine*. 2009;27:5488-95. doi: 10.1016/j.vaccine.2009.06.103. PubMed PMID: 19616494; PubMed Central PMCID: PMCPMC2745558.
- 187 Libraty DH, Zhang L, Woda M, Acosta LP, Obcena A, Brion JD, et al. Neonatal BCG vaccination is associated with enhanced T-helper 1 immune responses to heterologous infant vaccines. *Trials Vaccinol*. 2014;3:1-5. doi: 10.1016/j.trivac.2013.11.004. PubMed PMID: 24611083; PubMed Central PMCID: PMCPMC3943168.
- 188 Hatherill M, Cobelens F. Infant BCG vaccination is beneficial, but not sufficient. *Lancet Glob Health*. 2022;10:e1220-e1. doi: 10.1016/S2214-109X(22)00325-4. PubMed PMID: 35961334.
- 189 Hussey GD, Watkins ML, Goddard EA, Gottschalk S, Hughes EJ, Itoni K, et al. Neonatal mycobacterial specific cytotoxic T-lymphocyte and cytokine profiles in response to distinct BCG vaccination strategies. *Immunology*. 2002;105:314-24. doi: 10.1046/j.1365-2567.2002.01366.x. PubMed PMID: 11918693; PubMed Central PMCID: PMCPMC1782661.