



## Immunohistochemical Profiling of CD68 and VEGF in First-Trimester Miscarriage Placentas

Ghanyia Jasim Shanyoor<sup>1\*</sup>, Mukhtar Khamis Haba<sup>1</sup>, Ban Hadi Hameed<sup>2</sup>

<sup>1</sup>Department of Biology, College of Science for Women, University of Baghdad, Baghdad, Iraq; <sup>2</sup>Department of Obstetrics and Gynecology, College of Medicine, University of Mustansiriyah, Baghdad, Iraq

### ABSTRACT

**Background:** Miscarriage occurs when a pregnancy ends spontaneously before the fetus is viable outside the uterus. Early miscarriage, sometimes referred to as first-trimester miscarriage, which happens before 12 weeks. Vascular endothelial growth factor (VEGF), an angiogenic protein necessary for blood vessel formation, and cluster of differentiation 68 (CD68), a marker for macrophages, are vital to the immunological response. This research aimed to evaluate VEGF and CD68 expression patterns in first-trimester miscarriage in placental tissues and decidua in a bid to ascertain their potential function in immune response and angiogenesis in miscarriage.

**Methods:** The study included 60 women who experienced spontaneous abortion during the first trimester, 30 with missed abortion, and 30 with incomplete abortion. Using immunohistochemical staining, the expression of CD68 and VEGF in decidual and placental tissues was investigated. Expression levels were scored semi-quantitatively by averaging data from five fields per slide and subjected to statistical analysis.

**Results:** Building on these methods, immunohistochemical staining confirmed significantly elevated CD68 expression in both placental and decidual tissues from missed and incomplete abortions, indicating enhanced macrophage infiltration. Widespread staining (>50%) was seen in 50% of incomplete abortion decidua and 30% of missed abortion tissues. In contrast, VEGF expression was found to be predominantly negative in all samples, with no case showing positive staining, indicating impaired angiogenesis.

**Conclusion:** The findings highlight a dual pathological mechanism in early miscarriage characterized by enhanced inflammatory macrophage infiltration and deficient angiogenic support. This imbalance may contribute to placental dysfunction and pregnancy failure. These results underscore the potential diagnostic and therapeutic relevance of immune–angiogenic pathways in early miscarriage

**Keywords:** CD68; Immunohistochemical staining; Miscarriage; VEGF

*\*Corresponding author:*  
Ghanyia Jasim Shanyoor,  
Department of Biology,  
College of Science for Women,  
University of Baghdad,  
Baghdad, Iraq  
**Email:** ghaniya.idam2202p@  
csw.uobaghdad.edu.iq

*Cite this article as:*  
Shanyoor GJ, Haba MK, Hameed  
BH. Immunohistochemical  
Profiling of CD68 and VEGF  
in First-Trimester Miscarriage  
Placentas. *Iran J Immunol.* 2026;  
23(1): 34-43,  
doi: 10.22034/iji.2026.109210.3127.

Received: 2025-10-14  
Revised: 2025-12-24  
Accepted: 2026-02-09

## INTRODUCTION

Miscarriage is defined as a spontaneous and unintentional loss of a pregnancy before the fetus is viable. It replaced the term “abortion” as the medical definition of miscarriage (1, 2). Miscarriage before the twentieth week of pregnancy is caused by chromosomal abnormalities (3) or uterine or cervical disorders (4). Infections and other medical conditions account for approximately 15% of early and 66% of late miscarriages. Immunological and environmental factors, maternal age, chronic health conditions, and lifestyle factors like smoking, alcohol consumption, obesity, and stress can increase the risk of miscarriage, particularly in mothers aged 35 and older (5, 6). Miscarriage typically occurs in the first three months of pregnancy, with common symptoms including vaginal spotting and passing blood clots (7). Decidua is differentiated into three regions that include the decidua parietalis, decidua capsularis, and decidua basalis, in accordance with the assertion of Yin et al. 2024 (8). The placenta, a crucial fetal organ, rapidly develops during pregnancy, secreting signaling molecules that govern maternal-fetal interactions, influencing fetal development, growth, and immunity (9, 10). Additionally, the placenta plays a central role in providing nutrients, disposing of metabolic wastes, and ensuring a healthy pregnancy by facilitating complex communication between the fetus and mother, involving immune-related tasks (11). The interaction between the fetus and the maternal immune system is crucial for protecting against immune rejection. Factors like immune tolerance, placental macrophages, and cytokine release have been reported to be involved in miscarriages (12-14). The delicate balance between immune tolerance and immune responses to infections in a pregnancy is crucial for a healthy pregnancy, and any condition disturbing this balance can lead to miscarriage.

Cluster of differentiation 68 (CD68), a receptor and soluble protein, is a promising

candidate for therapeutic purposes (15). CD68 is a marker of macrophage presence and activation in tissues and helps diagnose immune reactions that accompany pregnancy or disease-associated miscarriages (Ref). Vascular endothelial growth factor (VEGF), a cytoprotective and anti-apoptotic factor, is essential in the formation of blood vessels in the placenta, ensuring adequate blood flow to the developing fetus. It plays a role in modifying the immune environment, protecting the fetus from mother’s immune responses. Abnormal VEGF levels can lead to pregnancy complications including miscarriage (16-20). Immunopathological mechanisms in miscarriage include dysregulated cytokine signaling, altered macrophage activation, and defective angiogenesis. Pro-inflammatory cytokines such as interleukin- IL)-18 and IL-23 can shift the immune environment toward a Th1-dominant response, promoting tissue damage and trophoblast apoptosis (21, 22). Concurrently, aberrant expression of angiogenic factors such as VEGF compromises placental vascularization, and altered macrophage activity, reflected by CD68 expression, contributes to inflammatory infiltration and impaired tissue remodeling (23).

The present study aimed to investigate the immunohistochemical expression patterns of CD68 and VEGF in placental tissues and decidua from first-trimester miscarriages to elucidate their potential roles in the pathophysiology of miscarriage.

## MATERIALS AND METHODS

### *Study Population*

A total of sixty women attending the Fatima Al Zahra Hospital and Al-Liqa in Baghdad, Iraq, were recruited. The participants included thirty women with missed abortion and thirty with incomplete abortion. These women experienced abdominal pain and vaginal bleeding in the first trimester of their pregnancies, which led to their hospitalization.

### *Inclusion and Exclusion Criteria*

The study included women aged 18-40 years who presented with Rh-positive blood type and no known cause of vaginal bleeding, while the exclusion criteria included samples of women who smoked or consumed alcohol, were receiving ongoing treatment for chronic diseases, had anemia (hemoglobin <8 g/dL), and had a negative Rh blood group. As women were brought for uterine evacuation at the women's emergency department of the hospital the products of conception were taken from the patients.

### *Ethical Approval*

Ethical approval for the present study was obtained from the Ethics Committee of the Baghdad University's College of Women's Science (Approval number: 7902/22) on December 21, 2023. The study was approved by the scientific committee of Bagdad University, college of science (no. 7902-22 on 21/12/2023). Their medical histories were recorded, and clinical examinations were conducted before enrollment. Participants were then assigned to different groups.

### *Immunohistochemical Analysis of CD68 and VEGF*

Curettage was used to create placenta specimens, which were embedded in paraffin blocks, preserved in 10% formalin, sliced into 5 µm pieces, deparaffinized in xylene, rehydrated, and incubated with 3% hydrogen peroxide at 37°C for 10 minutes to inhibit peroxidase activity. Slides were washed twice in phosphate buffer saline, rinsed with distilled water for 5 minutes, and incubated with primary antibodies against CD68 and VEGF. Slides were then washed three times in PBS, then rinsed twice in PBS and distilled water. The study involved applying mouse anti-CD68 and mouse anti-VEGF conjugates (Dako, Denmark), incubating at 37°C, and performing four PBS rinses. The mixture was then applied to tissue, rinsed twice, and counterstained with Harris Hematoxylin. The slides were then dehydrated, mounted, and

examined under a light microscope (24).

### *Assessment of the Immunohistochemical Staining*

Two experienced pathologists evaluated slides separately using a semi-quantitative grading program. Based on their cytoplasmic staining, placenta, and decidual stromal cells were categorized as negative (0), slightly positive (+1), moderately positive (+2), or strongly positive (+3). A noticeable brown stain receives a positive score. When more than 50% of the cells showed strong staining, the result was recorded as strong positive. Additionally, when less than 20% of the cells showed significant staining, the result was reported as slightly positive. When 20% to 50% of the target cells showed strong staining, the staining result was moderately positive (+2) (25).

### *Statistical Analysis*

At the end of the study, all data was compiled and tabulated. Appropriate tests were used for statistical analysis, such as Fisher's test for the immunohistochemical data and the Pearson chi-square (free statistics software).

## **RESULTS**

### *Demographic Data*

The mean age of the participants was 27.13±5.4 years in the missed abortion group, and 27.83±6.5 years in the incomplete abortion group with no significant differences. Additionally, there was no significant differences in pregnancy status and parity between the two groups, Table 1.

### *Expression of CD68 and VEGF in Placental and Decidual Tissues*

The immunohistochemical analysis revealed that the highest expression of CD68 in incomplete abortion cases was observed in the decidua, with strong expression (+++, >50%) detected in 15 out of 30 samples (50%).

**Table 1: The Number and percentage of Pregnancy status, children, and maternal among women participating in the study**

Variables	Missed abortion		Incomplete abortion		P-value	
	Frequency No. (30)	%	Frequency No. (30)	%		
Pregnancy status	1-2	9	30%	5	16.6%	0.24447 NS
	3-4	12	40%	10	33.3%	0.6317 NS
	≥5	9	30%	15	50.0%	0.2190 NS
Children	0-1	10	33.3	9	30.0%	0.8146 NS
	2-3	15	50%	10	33.3%	0.2354 NS
	>3	5	16.6	11	36.3%	0.1038 NS
Maternal age (years)	Mean±SD	27.13±5.4		27.83±6.5		0.1NS

**Table 2. The intensity of CD68 expression in decidual and placental tissues**

Grade (intensity of CD68)	Decidua (incomplete abortion; (n = 18)	Placenta tissue (incomplete abortion (n = 12)	Decidua (missed abortion; (n = 12)	Placenta tissue (missed abortion; (n = 18)	Statistical test
Negative (0) <10	0 (0%)	0 (0%)	0 (0%)	0 (0%)	
Weak (+; 10-20%)	0 (0%)	0 (0%)	0 (0%)	0 (0%)	
Moderate (++; 20-50%)	3 (10%)	12 (40%)	3 (10%)	9 (30%)	X <sup>2</sup> =0.75 (NS)
Strong (+++; >50%)	15 (50%)	0 (0%)	9 (30%)	9 (30%)	Fisher test= 0.78 (NS)
Total	18 (60)	12 (40)	12 (40)	18 (60)	0.12 (NS)

CD68: Cluster of differentiation; X<sup>2</sup>: Chi-square; NS: non-significant; \*p-value<0.05: significant; \*\*p-value < 0.01: highly significant


In contrast, 12 samples (40%) exhibited moderate expression (++ staining, 20–50%) in the placental tissue, as shown in Table 2. Meanwhile, the missed abortion group exhibited strong expression (+++, >50%) in both decidual and placental tissues, with 9 out of 30 samples (30%) in each region. Additionally, moderate expression (++ staining, 20–50%) was observed in the placental area of 9 out of 30 samples (30%). In cases of missed and incomplete abortion, moderate expression (++ staining, 20–50%) was less frequent in the decidua, occurring in only 10% of the samples.

In cases of incomplete abortion, 60% of the decidual samples showed negative (0) VEGF expression, while 40% of placental tissues in the same group exhibited negative VEGF expression. For missed abortion


cases, 40% and 60% of the decidual and placental tissues displayed negative VEGF staining, respectively. None of the samples exhibited weak (0-20%), moderate (20-50%), or strong (>50%) positive VEGF staining. In total, 60% of decidual samples from incomplete abortion cases and 60% of placental tissues from missed abortion cases showed negative VEGF expression. Similarly, 40% of both decidual and placental samples from missed abortion cases exhibited negative staining.

#### *Immunohistochemical Evaluation of Placental and Decidual Tissues*

Fig. 1 presents the immunohistochemical expression of CD68 across different study groups. Fig. 1A shows moderate CD68 expression in the placental tissue from an incomplete



**Fig. 1.** Immunohistochemical expression of CD68 in placental and decidual tissues. Representative micrographs showing moderate CD68 expression in: (A) placental tissue from an incomplete abortion, (B) placental tissue from a missed abortion, (C) decidual tissue from an incomplete abortion, and (D) decidual tissue from a missed abortion ( $\times 40$ ). CD68: Cluster of Differentiation 68.



**Fig. 2.** Immunohistochemical expression of CD68 in placental and decidual tissues. Representative micrographs showing strong CD68 immunoreactivity in: (A) placental tissue from an incomplete abortion, (B) placental tissue from a missed abortion, (C) decidual tissue from an incomplete abortion, and (D) decidual tissue from a missed abortion ( $\times 40$ ). CD68: Cluster of Differentiation 68.


abortion case. In Fig. 1B, moderate CD68 expression was observed in placental tissue from a missed abortion case. Fig. 1C illustrates moderate CD68 expression in the decidua of an incomplete abortion case. Finally, the decidua in a missed abortion case has moderate expression of CD68, as shown in Fig. 1D. The tissue of incomplete abortions expressed high levels of CD68 (Fig. 2A). Similarly, Fig. 2B displays strong expression of CD68 in the placental tissue of missed abortion cases. In Fig. 2C, strong CD68 expressions were observed in the decidua of incomplete abortion cases, while Fig. 2D shows strong expressions in the decidua of missed abortion cases. Fig. 3 shows the immunohistochemical expression of VEGF in the study groups. Fig. (3A) displays negative VEGF expression in the placental tissue of incomplete abortion cases. Similarly, Fig. (3B) shows negative expression in the placental tissue of missed abortion cases. A negative VEGF expression was observed in the decidua of incomplete and missed abortion cases, as shown in Figs. 3C and D.

## DISCUSSION

A previous study by Zhang and colleagues (2022) did not demonstrate a linear relationship between the risk of spontaneous abortion and maternal age (appeared J-shaped relationship). Additionally, this study showed that the optimal threshold for maternal age is around 29 years, and maternal age over 30 years is accompanied with increased risk of spontaneous abortion (26). An increase in Spontaneous abortion is strongly correlated with age and parity and both the number of children and the frequency of abortion increase in grand multiparas (27).

The present study provides insight into the immunological and angiogenic alterations associated with first-trimester miscarriage by examining CD68 and VEGF expressions in decidual and placental tissues from incomplete and missed abortions. The findings highlight a dysregulated immune–vascular environment that may contribute to miscarriage.

The marked CD68 expression observed in



**Fig. 3.** Immunohistochemical expression of VEGF in placental and decidual tissues. Representative micrographs showing negative VEGF immunoreactivity in: (A) placental tissue from an incomplete abortion, (B) placental tissue from a missed abortion, (C) decidual tissue from an incomplete abortion, and (D) decidual tissue from a missed abortion ( $\times 40$ ). VEGF: Vascular Endothelial Growth Factor.

both decidual, and placental tissues indicates enhanced macrophage activity, supporting the concept that immune imbalance plays a central role in miscarriage. Although macrophages are essential for implantation, tissue remodeling, and maintenance of maternal–fetal immune tolerance, excessive accumulation or polarization toward a pro-inflammatory (M1) phenotype can be detrimental (28, 29). The high proportion of samples showing strong CD68 expression in the present study suggests that uncontrolled macrophage activation may contribute to tissue damage, trophoblast apoptosis, and impaired placental function.

In contrast, VEGF expression was largely absent in both decidual and placental tissues, indicating compromised angiogenic signaling. VEGF plays a critical role in placental vascular development and fetal growth, and its downregulation has been associated with placental insufficiency and adverse pregnancy outcomes (30, 31). The concurrent high CD68 expression and low VEGF levels supports the concept that excessive immune activation may suppress angiogenic pathways, leading to inadequate placental vascularization. This imbalance between inflammation and angiogenesis may represent a key mechanism underlying miscarriage. The histological evidence of increased CD68-positive macrophages suggests an inflammatory environment in which immune cells may clear apoptotic debris or respond to tissue injury, but cause further placental damage (32). Simultaneously, reduced VEGF expression indicates impaired vascular support, reinforcing the concept that failed pregnancies may result from a shift from immune tolerance and angiogenic support toward inflammation and vascular insufficiency.

Previous studies support the present findings that immune dysregulation and impaired angiogenesis play key roles in early miscarriage. Increased infiltration of CD68-positive macrophages has been consistently reported in decidual and placental tissues

from spontaneous and missed abortions, suggesting an exaggerated inflammatory response that disrupts trophoblast function and maternal–fetal tolerance (33–35). In parallel, reduced VEGF expression has been associated with defective placental vascular development and poor pregnancy outcome (36, 37). Additionally, inflammatory cytokine activity has been shown to suppress angiogenic signaling, linking immune activation to vascular insufficiency (38). These findings support our observation that increased macrophage activity combined with reduced VEGF expression contributes to the pathogenesis of first-trimester miscarriage.

#### *Study Limitation*

Finally, this study is limited by its small sample size, restriction to Rh-positive women, and lack of healthy first-trimester control placentas. These factors limit the generalizability of the findings and prevent definitive comparisons with normal placental development.

Semi-quantitative immunohistochemical scoring was used to evaluate CD68 and VEGF expressions because this method allows simultaneous assessment of both staining intensity and tissue localization within placental and decidual structures. This approach is widely used in histopathological studies when spatial distribution of immune cells and angiogenic markers is relevant. Although digital image analysis and molecular techniques such as RT-PCR or ELISA provide more objective quantification, they do not preserve tissue architecture or allow cell-specific localization. Nevertheless, we acknowledge that the lack of molecular quantification represents a limitation, and future investigations will incorporate digital image analysis and complementary molecular assays to strengthen quantitative validation of these findings. We acknowledge that the use of only chi-square and Fisher's exact tests without adjustment for multiple comparisons or confounding factors represents another

limitation in our study. Due to the limited sample size and the categorical nature of the immunohistochemical data, more advanced statistical modeling was not feasible. Future studies with larger cohorts will include multivariate analyses and appropriate correction methods (e.g., Bonferroni or false discovery rate adjustments) to improve statistical rigor

## CONCLUSION

The present study reveals that first-trimester miscarriage placentas, from both incomplete and missed abortions, have a unique immunohistochemical profile. They feature high CD68 and low or absent VEGF expressions. The strong CD68 staining in decidual tissues indicates an active inflammatory process. In contrast, the near-total lack of VEGF expression in all tissue samples points to a serious impairment in angiogenesis, which is crucial for placental development and fetal support. The concurrent increased CD68 and decreased VEGF expressions suggests an environment where immune activation coincides with a lack of vascular support. These findings highlight the roles of immune dysregulation and impaired angiogenesis in miscarriage; however, the limited sample size and cross-sectional design restrict causal interpretation. Future studies incorporating larger cohorts and functional analyses are needed to clarify underlying mechanisms and therapeutic potential.

## ACKNOWLEDGEMENTS

The authors wish to thank the management of the Department of Biology, College of Science for Women, University of Baghdad, Baghdad, Iraq for granting the permission to use the departmental facilities and thank to the Fatima Al Zahra Hospital in Baghdad, Iraq for their help recruit pregnant women.

## CONFLICTS OF INTEREST

The authors declare no conflicts of interest.

## REFERENCES

1. Rouse Cynthia E., Eckert Laura O., Babarinsa Idowu, Fay Elizabeth, Gupta Manisha, Harrison Mary S., et al. Spontaneous abortion and ectopic pregnancy: Case definition & guidelines for data collection, analysis, and presentation of maternal immunization safety data. *Vaccine*. 2017;35(48 Pt A):6563-74.
2. Malory B. The transition from abortion to miscarriage to describe early pregnancy loss in British medical journals: a prescribed or natural lexical change? *Med Humanit*. 2022;48(4):489-96.
3. Akram GT, Kandala NJ. Association between *Leptotrichia amnionii* and *Atopobium vaginae* as a risk factor for miscarriage states in specimens of Iraqi women. *Iraqi Journal of Science*. 2021;62(1):83-95.
4. Magnus MC, Morken NH, Wensaas KA, Wilcox AJ, Håberg SE. Risk of miscarriage in women with chronic diseases in Norway: A registry linkage study. *PLoS Med*. 2021;18(5):e1003603.
5. La X, Wang W, Zhang M, Liang L. Definition and Multiple Factors of Recurrent Spontaneous Abortion. *Adv Exp Med Biol*. 2021;1300:231-57.
6. Zhang M, Yang BY, Sun Y, Qian Z, Xaverius PK, Aaron HE, et al. Non-linear relationship of maternal age with risk of spontaneous abortion: a case-control study in the China birth cohort. *Front Public Health*. 2022;10:933654.
7. Turkey EF, Hamad AG, AL-Mansoori AK. (2022): The association of prothrombin gene mutations and cytomegalovirus infection with abortion among Iraqi women. *Baghdad Sci J*. 2022;19(4):768-74.
8. Yin Z, Su J, Lu L, Yang L, Su S, Jiang X. Visual identification of three kinds of human decidual tissues from elective termination of pregnancy. *Placenta*. 2024;146:89-100.
9. Burton GJ, Fowden AL. The placenta: a multifaceted, transient organ. *Philos Trans R Soc Lond B Biol Sci*. 2015;370(1663):20140066.
10. Rasheed YM. Applying B-mode ultrasound for determining pregnancy by means of measurement of placentome diameter and fetal Bi-parietal diameter in dairy cows. *Iraqi J Vet Med*. 2018;42(2):90-6.
11. Estrela D, Santos RF, Masserdotti A, Silini A, Parolini O, Pinto IM, et al. Molecular Biomarkers for Timely and Personalized Prediction of Maternal-Fetal Health Risk. *Biomolecules*.

- 2025;15(3):312.
12. Hussain T, Murtaza G, Kalhoro DH, Kalhoro MS, Yin Y, Chughtai MI, et al. Understanding the Immune System in Fetal Protection and Maternal Infections during Pregnancy. *J Immunol Res.* 2022;2022:7567708.
  13. Moldenhauer LM, Hull ML, Foyle KL, McCormack CD, Robertson SA. Immune-Metabolic Interactions and T Cell Tolerance in Pregnancy. *J Immunol.* 2022;209(8):1426-36.
  14. Ahmad ZA, Idan EM, Ardalan NM. Role of IL-37 and dectin-1 during toxoplasmosis. *Baghdad Sci J.* 2023;20(3):746-54.
  15. Daub K, Siegel-Axel D, Schönberger T, Leder C, Seizer P, Müller K, et al. Inhibition of foam cell formation using a soluble CD68-Fc fusion protein. *J Mol Med (Berl).* 2010;88(9):909-20.
  16. Jiang X, Wang J, Deng X, Xiong F, Zhang S, Gong Z. The role of microenvironment in tumor angiogenesis. *J Exp Clin Cancer Res.* 2020;39(1):204.
  17. Khalil KH, Al-Hassawi B, Abdo J. Correlation of neuroendocrine differentiation with neuroendocrine cell hyperplasia and vascular endothelial growth factor in colorectal
  18. Qaddoori YB, Al-Khafaji AS, Khashman BM, Ghafour KHA. The study of histone deacetylase immunosuppression in relation to regulating vascular endothelial growth factor (VEGF) has implicated it in the malignant progression of colorectal cancer. *Iraqi Journal of Science.* 2025;66(5):1861-1875.
  19. Gualdoni GS, Jacobo PV, Barril C, Ventureira MR, Cebral E. Early Abnormal Placentation and Evidence of Vascular Endothelial Growth Factor System Dysregulation at the Feto-Maternal Interface After Periconceptional Alcohol Consumption. *Front Physiol.* 2022;12:815760.
  20. Hussein A. H. (2024): Studying the effect of thymol-containing nanoemulsion extracted from thyme on angiogenesis-controlling genes VEGF and VEGFR in a human liver cancer cell line. *Baghdad Sci J.* 2025;22(4):1151–62
  21. Löb S, Ochmann B, Ma Z, Vilsmaier T, Kuhn C, Schmoedel E, et al. The role of interleukin-18 in recurrent early pregnancy loss. *J Reprod Immunol.* 2021;148:103432.
  22. Zhao QY, Li QH, Fu YY, Ren CE, Jiang AF, Meng YH. Decidual macrophages in recurrent spontaneous abortion. *Front Immunol.* 2022;13:994888. doi:10.3389/fimmu.2022.994888.
  23. Andraweera PH, Dekker GA, Roberts CT. The vascular endothelial growth factor family in adverse pregnancy outcomes. *Hum Reprod Update.* 2012;18(4):436–457. doi:10.1093/humupd/dms011.
  24. Hilmi MN, Mirza SA, Al-Jaleeli AN. The Sensitivity of Immunohistochemical Expression of p53 as an Indicator of the Malignant Potential of Gastric Hyperplastic Polyps: A Retrospective Study. *Al-Rafidain J Med Sci.* 2024;7(1):198-202.
  25. Lv S, Yang Y, Kwon S, Han M, Zhao F, Kang H, et al. The association of CXCR4 expression with prognosis and clinicopathological indicators in colorectal carcinoma patients: a meta-analysis. *Histopathology.* 2014;64(5):701-12.
  26. Zhang, M., Yang, B. Y., Sun, Y., Qian, Z., Xaverius, P. K., Aaron, H. E., ... & Yue, W. (2022). Non-linear relationship of maternal age with risk of spontaneous abortion: a case-control study in the China birth cohort. *Frontiers in Public Health,* 10, 933654
  27. Cohain JS, Buxbaum RE, Mankuta D. Spontaneous first-trimester miscarriage rates per woman among parous women with 1 or more pregnancies of 24 weeks or more. *BMC Pregnancy Childbirth.* 2017 Dec;17(1):437. <https://doi.org/10.1186/s12884-017-1620-1>.
  28. Mantovani A, Biswas SK, Galdiero MR, Sica A, Locati M. Macrophage plasticity and polarization in tissue repair and remodelling. *J Pathol.* 2013;229(2):176-85.
  29. Tsao FY, Wu MY, Chang YL, Wu CT, Ho HN. M1 macrophages decrease in the decidua from normal pregnancies but not from spontaneous abortions or unexplained recurrent spontaneous abortions. *J Formos Med Assoc.* 2018;117(3):204-211.
  30. Vuorela P, Hatva E, Lymboussaki A, Kaipainen A, Joukov V, Persico MG, et al. Expression of vascular endothelial growth factor and placenta growth factor in the human placenta. *Biol Reprod.* 1997;56(2):489-94.
  31. Zygmunt M, Herr F, Münstedt K, Lang U, Liang OD. Angiogenesis and vasculogenesis in pregnancy. *Eur J Obstet Gynecol Reprod Biol.* 2003;110 Suppl 1:S10-8.
  32. Wu HM, Chen LH, Hsu LT, Lai CH. Immune Tolerance of Embryo Implantation and Pregnancy: The Role of Human Decidual Stromal Cell- and Embryonic-Derived Extracellular Vesicles. *Int J Mol Sci.* 2022 2;23(21):13382.
  33. Bulmer JN, Morrison L, Longfellow M, Ritson A, Pace D. Granulated lymphocytes in human endometrium: histochemical and immunohistological studies. *Placenta.* 1991;12(3):295–304
  34. Liu S, Diao L, Huang C, Li Y, Zeng Y, Kwak-Kim J. The role of decidual immune cells on human pregnancy. *J Reprod Immunol.* 2013;97(1):1–10.
  35. Zhou Y, Fu B, Xu X, Zhang J, Tong X, Wang Y. Aberrant macrophage polarization in the decidua of women with recurrent spontaneous abortion. *Reprod Biol Endocrinol.* 2018;16:1–10.
  36. Charnock-Jones DS, Kaufmann P, Mayhew TM.

- Aspects of human fetoplacental vasculogenesis and angiogenesis. *Placenta*. 2004;25(2-3):103-113.
37. Liu Y, Fan X, Wang R, Lu X, Dang YL, Wang H. Abnormal expression of VEGF and its receptors in placental tissues from spontaneous abortion. *Mol Med Rep*. 2015;12(3):399-404.
38. Wang WJ, Hao CF, Yi L, Yin GJ, Bao SH, Qiu LH, et al. Increased prevalence of T helper 17 cells in decidua and peripheral blood in unexplained recurrent spontaneous abortion. *Am J Reprod Immunol*. 2014;72(5):486-495.