

REVIEW ARTICLE

The Impact of Food Vitamins and Minerals on Proliferation, Differentiation, Homeostasis and Regenerative Capacity of Body Mesenchymal Stem Cells: A Narrative Review

Somayeh Montazeri¹, Marziyeh Hamzavi^{2*}, Mehra Nazempour³, Atefeh Sadat Masoumi⁴, Davood Mehrabani^{5,6,7*}, Seyed Jalil Masoumi⁸

1. Comparative and Experimental Medicine Center, Shiraz University of Medical Sciences, Shiraz, Iran

2. Hamzavi Dental Clinic, Shiraz, Iran

3. Nutrition Research Center, Department of Clinical Nutrition, School of Nutrition and Food Sciences, Shiraz University of Medical Sciences, Shiraz, Iran

4. School of Dentistry, Islamic Azad University, Shiraz Branch, Shiraz, Iran

5. Medical Imaging Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

6. Stem Cell Technology Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

7. Burn and Wound Healing Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

8. Department of Biomedical and Tissue Engineering, Science and Research Branch, Islamic Azad University, Tehran, Iran

ARTICLE INFO

Keywords:

Food

Diet

Vitamins

Minerals

Mesenchymal stem cells

ABSTRACT

Several lines of evidence confirmed the idea that stem cells are among the major players in orchestrating the response of our body to nutrients, mainly due to their crucial role in tissue homeostasis as dietary components and nutritional status can significantly impact stem cell survival and performance in human body and subsequently health status. This review concentrated on literature findings to determine the impact of food components on function, proliferation, differentiation, homeostasis, and regenerative capacity of body mesenchymal stem cells by searching related databases and focusing on related keywords from June 2000 to June 2026. Even many studies regarding the impact of vitamins and mineral components on proliferation, differentiation, homeostasis and regenerative capacity of body mesenchymal stem cells are *in vitro* and require *in vivo* validation; but the positive physiological effects of vitamins and minerals through several mechanisms in the literature reflect their pivotal role in regenerative medicine. It seems that improving nutritional intake and biochemical status of key micronutrients (e.g. vitamins and minerals) can be supportive for the efficacy of MSCs in therapies and immune function and subsequently the recovery and clinical outcome of patients. A proper food intake can particularly target the enhancement of stem cell activity, epigenetic regulation, redox-inflammatory pathways and cellular metabolism. In this review, mitochondrial metabolic checkpoint regulating stem cell maintenance was highlighted. It was discussed how diet affects the metabolism of stem cells through the nutrient sensing pathways, thereby affecting stem cell fate decision and tissue regeneration.

*Corresponding author:

Marzieh Hamzavi, Davood

Mehrabani, PhD;

Medical Imaging Research Center,
Shiraz University of Medical
Sciences, Shiraz, Iran.

Email: mahshamzavi@gmail.com

davood_mehrabani@yahoo.com

Received: March 1, 2026

Revised: May 25, 2026

Accepted: June 1, 2026

Please cite this article as: Montazeri S, Hamzavi M, Nazempour M, Masoumi AS, Mehrabani D, Masoumi SJ. The Impact of Food Vitamins and Minerals on Proliferation, Differentiation, Homeostasis and Regenerative Capacity of Body Mesenchymal Stem Cells: A Narrative Review. Int J Nutr Sci. 2026;11(3):1-18. doi:

Introduction

Nutrients can directly affect stem cells or indirectly via regulating the stem cell niche. Also, nutrients can regulate hormone production, which in turn can impact the behavior of stem cells and their niche (1). Several lines of evidence confirmed the idea that stem cells are among the major players in orchestrating the response of our body to nutrients, mainly due to their crucial role in tissue homeostasis as dietary components and nutritional status can significantly impact stem cell survival and performance in human body and subsequently health status (2-7). In doing so, tissue stem cells not only use nutrients for their metabolic needs but also adapt their activities, such as self-renewal, autophagy or differentiation to the metabolic environment and nutrient availability (1). Micronutrients and dietary components are among the major signaling molecules that not only influence the cell metabolism; but also can modulate stem cell activity. In particular; minerals, vitamins, etc. are important to maintain stem cell proliferation and self-renewal via signaling pathways and gene expression networks and epigenetic regulation. So the regulation in nutrition can be an emerging target to control stem cell function, survival and pools in body tissues (8-10). In this relation, the concept of functional food components is crucial to provide a physiological and health benefits relevant to stem cell function (11). The functional food micronutrients can potentially enhance regenerative outcomes and protect stem cells from stress and to be novel perspectives for health promotion and disease prevention (12).

Among the functional food micronutrients, vitamins and minerals are provided to human body from natural sources of vegetables, fruits, oils, nuts, meat, dairy products, etc. So selected micronutrients found in foods can have specific effects at cellular level (13). It was shown that functional food components have various physiological advantages in human body as plausible modulators of stem cell function and enhancing regenerative outcomes in several diseases (3, 14).

Food-gene interaction was depicted as a leading risk factor for some diseases like inflammatory bowel disease (IBD) and colorectal cancer (CRC). Recent studies in stem cell biology demonstrated that diet and nutrition by signaling to intestinal MSCs (I-MSCs) by shifting nutrient-sensing transcriptional functions could impact barrier integrity and susceptibility to inflammation and tumorigenesis (15). Some nutrient ingredients and dietetic regimens were shown to have beneficial effects on the regulation of stem cells and tissue homeostasis and can lead to inhibition of

the cancer transformation process and progression and prevention of the onset of a drug resistance (16). Nutrients can normally affect the physiology of stem cells due to capability of several nutrient derived metabolites, released during the catabolic process, to induce chromatin reshaping, epigenetic modifications and gene expression modulation (17).

Indeed, nutrients may play a direct or indirect role by regulating the stem cell niche and can regulate hormone production, which in turn can affect the behavior of stem cells and their niche. In response to these stimuli, stem cells can activate signaling pathways, reprogram their metabolism and gene expression, and convert the dietary input into fate decisions. For stem cells, the main molecular node connecting diet and function was depicted by the adenosine monophosphate-activated protein kinase (AMPK)-mammalian target of rapamycin (mTOR)-sirtuin 1, (SIRT1) pathway (1). Even though these pathways are common to stem cells, the impact of diet on stem cells may be dramatic for various reasons. Stem cells have unique metabolic needs, which can be different based on their developmental stage (18-20).

Activation of these metabolic pathways is necessary for stem cell activities, inducing nutrient dependencies more profound than those in differentiated cells (21). Also, stem cells are characterized by lower levels of reactive oxygen species (ROS) when compared to their more differentiated counterparts; which are greatly affected by diet and nutrients and are important regulators of the balance between cell selfrenewal and differentiation (22). Therefore, this review has searched findings available in the literature to determine the impact of food components on body mesenchymal stem cells' function, proliferation, differentiation, homeostasis and regenerative capacity.

Materials and Methods

A comprehensive literature search was carried out utilizing databases of Scopus, Web of Science, PubMed, Cochrane, Science Direct, Ovid and Google Scholar by focusing on keywords of "food", "nutrition", "diet", "minerals", "vitamins", "mesenchymal stem cells", "cell function", "cell proliferation", "cell differentiation", "homeostasis", "regenerative capacity" and "healing" from June 2000 to June 2026.

Mesenchymal stem cells

Mesenchymal stem cells (MSCs) as multipotent adult stem cells having self-renewal and differentiation properties into specialized cell types

such as osteoblasts (bone cells), adipocytes (fat cells), and chondrocytes (cartilage cells) in order to maintain homeostasis and tissue regeneration in human being based on their anti-inflammatory and immunomodulatory characteristics and activities to repair damaged tissues (23-25). Their immunomodulatory characteristics can impact the activity of different immune cell types. They can promote polarization of macrophages from a pro-inflammatory (M1) to an anti-inflammatory (M2) phenotype, and enhance regulatory T cell populations making MSCs beneficial in conditions such as inflammation during tissue damages (26).

They secrete several bioactive factors such as cytokines and growth factors, which not only protect tissue regeneration and repair; but also modulate the local immune responses. This regenerative potential is enhanced by their involvement in the process of new blood vessel formation, which is crucial to supply injured tissues with nutrients and oxygen in order to promote wound healing by facilitating cellular communication and extracellular matrix (ECM) remodeling (27). They have the regenerative potential of migrating to sites of injury from distant organs and have the potential to exert paracrine effects *in situ* making them a proper candidate to counteract many diseases and degenerative conditions (28). MSCs have been dispersed throughout the body in various tissues such as bone marrow (29), adipose tissue (30), dental pulp (31), Wharton's jelly (32) endometrium (33) and menstrual blood (34). The therapeutic application of MSCs is intrinsically correlated to their potential for robust proliferation, effective self-renewal, and resistance to cellular senescence (3). They have been successfully utilized in treatment of many diseases such as infertility (35), spinal cord injury (36), burns (37), osteoarthritis (38), liver injuries (39), etc. MSCs have been labeled to track the cell fate after transplantation to confirm their share in healing process too (40).

Biological function of MSCs was shown to be modulated by environmental factors such as dietary components including minerals, vitamins, etc. which can affect cell proliferation, aging, inflammatory response and resistance to oxidative stress (3). Growing body of evidences revealed that minerals and vitamins are pivotal modulators in MSCs function, proliferation, differentiation, and homeostasis (5, 7, 41-45). The profound immunomodulatory capabilities of MSCs are crucial for their therapeutic efficacy in inflammatory and autoimmune diseases. These activities are not static; but are modulated by microenvironmental factors such as vitamins A, B3, B6, C, D, E and K together with minerals like magnesium, selenium,

zinc, iron and lithium chloride which are important for immunosuppressive and anti-inflammatory characteristics of MSCs because of a change in the profiles of cytokine secretion, surface marker expression, and intracellular signaling (3).

Vitamins

Vitamins are involved in various cellular activities, and their deficiency often leads to serious symptoms to people, sometimes even death. Most vitamins can be obtained through balanced food intake, and vitamin supplements are also extensively applied in healthcare practices. Different vitamins possess regulatory mechanisms at cellular level, particularly in stem cells. Functional stem cells need a culture system in which all components including water, inorganic salts, growth factors, amino acids, buffering reagent, energetic substrates, extracellular matrix, vitamins, vitamin-like organic factors are suitably balanced. So the regulation of nutrition is emerging as a viable target for stem cell modulation, which can impact not only cell survival; but also pluripotency and cell fates (41).

Human vitamins are generally divided into two groups of water-soluble vitamins and fat-soluble vitamins. Water-soluble vitamins are consisted of the B type vitamins and vitamin C, and fat-soluble vitamins compromise vitamins A, D, E and K. Some vitamins can be synthesized in the body, but at a very low rate; but all vitamins can be provided from food to fulfill the nutritional needs. Vitamins B1, B3, B6 and B7 participate in glucose metabolism of glycolysis, pentose pathway, glycogenolysis and gluconeogenesis; while vitamins B2, B3 and B5 are involved in fatty acid synthesis and degradation and vitamins B3, B6, B9 and B12 are essential in amino acid degradation. Vitamin C as an antioxidant suppresses generation of ROS. Vitamin A acts as an antioxidation and transcriptional regulation factor; vitamin D is a hormone that binds to nuclear receptors and regulates transcription, and is involved in calcium absorption; vitamin E is a strong fat-soluble antioxidant and modulates signal transduction and vitamin K is a cofactor for γ -glutamyl carboxylase that is crucial in blood clotting. (41). Six vitamins of B1, B2, B3, B5, B6 and B9 are necessary in cell proliferation. So in cell cultures, Basal Medium Eagle (BME), Minimum Essential Medium (MEM), Dulbecco's Modified MEM (DMEM) and DMEM, Ham's F12, RMPI, IMDM and α -MEM were developed with various vitamins (41).

The MSCs potentials to facilitate tissue repair is prominently affected by the local biochemical environment of different vitamins that can play a

direct role in modulating regenerative activities. It was shown that the differentiation of MSCs is effectively modulated by a wide range of vitamins that play a crucial role in tissue homeostasis and regenerative applications (3). In this relation, vitamins A, B, C, D, E and K have been highlighted as key micronutrients for stem cell activity (41, 46).

Vitamin A

Vitamin A is practically a group of compounds known as retinoids, including retinol, retinal and retinoic acid (RA). Vitamin A compounds are normally present in food of animal origin; while their precursor known as carotenoid is found in plants. Humans can synthesize vitamin A from carotenoids like β -carotenes that can be converted into two retinals that can be further reduced to retinol. Retinol either is then esterified into retinyl palmitate for storage, or is oxidized into RA. Vitamin A has also distinctive roles in embryogenesis and stem cells in culture as retinol and retinal are readily oxidized in culture and function as antioxidants to promote cell survival and growth (41).

RA, also known as tretinoin, is a bioactive derivative of vitamin A that mediates transcriptional programs critical for cell proliferation, differentiation and development (47). RA can enhance MSCs potential in suppression of T cell proliferation (48) and activate pro-inflammatory pathways within MSCs (49). RA *in vitro* with human umbilical cord-MSCs (UC-MSCs); *in vivo* in mouse model of liver diseases and *ex vivo* in liver inflammation co-culture model of human periodontal ligament-MSCs (PDL-MSCs) could enhance immunosuppression by suppressing proliferation of CD4⁺ and CD8⁺ T cells; reduction in programmed death-ligand-1 (PD-L1) expression; a significant rise in expression of human leukocyte antigens (HLA) class I and class II; a decrease in tumor necrosis factor-alpha (TNF- α) secretion and activating the pro-inflammatory nuclear factor kappa-light-chain-enhancer of activated B cells/nucleotide-binding-domain, leucine-rich repeat containing protein 3 (NF- κ B/NLRP3) axis and promoting the production of the proinflammatory cytokine of interleukin-1 beta (IL-1 β) (48, 49).

Vitamin A (retinol) and its metabolites (4-OH-RA and 4-Oxo-RA) roles in cell development were exhibited (50). In weaned piglets, dietary vitamin A could significantly affect I-MSC differentiation and stemness (51).

Vitamin B3

Vitamin B3 is a water-soluble B vitamin found naturally in many foods in common forms of niacin (nicotinic acid), nicotinamide (NAM)

and nicotinamide riboside (NR) as precursors of nicotinamide adenine dinucleotide (NAD) and nicotinamide adenine dinucleotide phosphate (NADP) that are cofactors or substrates in a wide range of metabolic reactions. Nicotinamide in medium can easily cross plasma membrane and enter the cytoplasm (41). Niacin acts in the body as a coenzyme, with more than 400 enzymes dependent on it for several reactions to help converting nutrients into energy, create cholesterol and fats, create and repair DNA, and exert antioxidant effects (52).

Vitamin B3 when treated with mouse adipose tissue derived-MSCs (Ad-MSCs) could regulate histone deacetylation and telomere maintenance via activating SIRT1, nicotinamide adenine dinucleotide (NAD⁺)-dependent histone deacetylase to postpone senescence and support differentiation (53). It was shown that nicotinamide and pyridoxine could stimulate muscle stem cell expansion and enhance regenerative capacity during aging as skeletal muscle is dependent on resident muscle MSCs (Mu-MSCs) for growth and repair. Nicotinamide and pyridoxine synergize through CK1-mediated cytoplasmic β -catenin activation and protein kinase B (AKT) signaling to promote amplification and differentiation of Mu-MSCs. Oral intake of nicotinamide and pyridoxine can accelerate muscle regeneration *in vivo* by stimulating Mu-MSCs, increase muscle strength and overcome Mu-MSC dysfunction and regenerative failure during aging (54).

Vitamin B6

Vitamin B6 is a water-soluble vitamin naturally found in several foods. Pyridoxal 5' phosphate (PLP) and pyridoxamine 5' phosphate (PMP) are the active coenzyme forms of vitamin B6. This vitamin is involved in gluconeogenesis and glycogenolysis, immune function, and hemoglobin formation (55). Administration of vitamin B6 was depicted to accelerate the clearance of transplanted MSCs from the body in a murine model that can be important in therapeutic applications (48). In murine model of hepatitis treated with UC-MSCs, an acceleration in clearance of MSCs from the body happened. It led to lower numbers of detectable UC-MSCs 4h post-injection, especially when combined with retinoic acid (48). Vitamin B6 can modulate the immunogenic and immunoregulatory profile of MSCs via a rise in the surface expression of HLA molecules; while upregulates PD-L1 immunomodulatory molecule and interleukin-1 receptor antagonist (IL-1RA) anti-inflammatory factor (48).

Vitamin C

Vitamin C, or l-ascorbic acid (AA/LAA), is a

water soluble vitamin due to its sugar-like structure. It is a strong antioxidant that reduces ROS that acts as a kinase inhibitor too. Its roles in promoting cell proliferation and survival, cell differentiation and reduction of cell senescence during reprogramming was demonstrated via suppressing p53 (41). It is an enzyme cofactor of critical importance in synthesis of collagen that can strongly support formation of ECM, which is necessary in tissue integrity, regeneration and remodeling (56, 57). It can improve neovascularization and tissue repair (58). In *in vivo* mice model of ischemic limb, transplantation of subchondral bone mesenchymal stem cells (SCB-MSCs) and AdMSCs was shown to enhance regeneration and tissue repair in bone defects acting as a cofactor in supporting ECM formation and deposition, collagen synthesis and improving blood perfusion and neovascularization (8, 56-66). Vitamin C contributes to immune regulation by suppressing the senescence-associated secretory phenotype (SASP) in MSCs and a decrease in production of proinflammatory cytokines, thereby helping maintain the functional integrity of MSCs and supporting the activity of immune effector cells (58, 67). Vitamin C as a potent antioxidant can directly scavenge ROS and protect mitochondrial homeostasis, and as a result can support MSCs from oxidative damage and the enhanced senescence via inhibiting ROS production through key signaling pathways (56, 63, 68-71). Vitamin C in presence of human bone marrow derived stem cells (BM-MSCs) and Ad-MSCs could induce osteogenesis in senescent MSCs and activate telomerase and expression of runtrelated transcription factor 2 (Runx2), alkaline phosphatase (ALP) and collagen type I alpha 1 chain (COL1A1) which are necessary in collagen synthesis and enhancing mineralization (56, 72, 73). Vitamin C when exposed to mouse embryonic mesoderm-derived MSCs could contribute to adipogenesis by a suppression of adipocytes differentiation (74). It was shown that vitamin C in addition to Glial-derived stem cells (GS-MCs) and Ad-MSCs resulted in chondrogenesis through enhancing differentiation and supporting chondrocytes from oxidative stress of hydrogen peroxide (H_2O_2) (7, 75). In *in vivo* mice model of ischemic limb, SCB-MSCs and Ad-MSCs in addition to vitamin C could enhance regeneration in bone defects and support ECM formation and tissue repair via acting as a cofactor in collagen synthesis and deposition and improving blood perfusion and neovascularization (8, 56-66). This vitamin with lung MSCs in rat model could promote vascular repair, bone regeneration and lung development because of modulating the immune environment; promoting the osteogenic differentiation; improving adhesion

and migration of endothelial progenitor cells (EPCs) and alleviating TNF- α -mediated inflammation (44, 76-78). Vitamin C when treated with SCS-MSCs were indicated to decrease proinflammatory phenotype of SASP, TNF- α , IL-1 β , IL-6 and IL-8 and support activation of natural killer (NK) cells and cytotoxic T lymphocytes (58). This vitamin *in vitro* when added to Ad-MSCs and lithium chloride increased cell proliferation and protected the cells from apoptosis via an increase in expression of the Bcl-2 gene and a significant reduction in Bax gene (14). Vitamin C with Ad-MSCs *in vivo* in mouse model of acute radiation syndrome (ARS) increased the survival rate following a lethal irradiation and reduced the proinflammatory cytokines, ECM formation, adhesion properties and boosted anti-inflammatory properties, detoxification and anti-oxidative stress control (79). This vitamin with Ad-MSCs demonstrated strong antioxidant activity and decreased ROS due to scavenging and inhibition of ROS production via protein kinase B/AKT/mTOR axis (56, 59, 63, 68, 80). Vitamin C in combination with gingival MSCs and mouse Ad-MSCs could directly modulate DNA and histone demethylation by participating as a cofactor for ten-eleven translocation (TET) enzymes, promoting an active DNA demethylation from 5-methylcytosine (5mC) to 5-hydroxymethylcytosine (5hmC); supporting Jumonji (JHDM) histone demethylases such as histone H3 lysine 36 dimethylation/trimethylation/2/3 (H3K36me2/3) and lead to an increase in cellular myelocytomatosis/Krüppel-like factor 4 (c-Myc/Klf4) and repression of protein 21 (p21) (7) (64) (81).

Vitamin D

Vitamin D (also named calciferol) is fat-soluble and biologically inert that is naturally present in a few foods and is produced endogenously when ultraviolet (UV) rays from sunlight is exposed to the skin and stimulate vitamin D synthesis. It must undergo two hydroxylations in liver and kidney for activation converting vitamin D to 25-hydroxyvitamin D [25(OH)D], also known as calcidiol 1,25-dihydroxyvitamin D [1,25(OH)2D], also known as calcitriol (82). The key role of the nutrient environment, and vitamin D signaling was illustrated in defining contribution of at least two different stem cell populations to mucosal homeostasis (83). Vitamin D was shown to directly enhance the regenerative capacity of MSCs, promote bone regeneration by modulating the local immune environment and directing osteogenic differentiation (44, 46). Also, it participates in angiogenesis and vascular repair by improving the endothelial progenitor cells activity (76, 78). In rat model, lung

MSCs were demonstrated to promote vascular repair, bone regeneration and lung development via modulation of the immune environment; promotion in osteogenic differentiation; improvement in endothelial progenitor cells (EPC) adhesion and migration; and alleviation of TNF- α -mediated inflammation (44, 76-78). Vitamin D has a potent immunomodulatory impact via the decrease in secretion of proinflammatory cytokines mediated by an inhibition in NF- κ B signaling (84-87). This mechanism can enhance the capability of MSCs in suppression of proliferation of T cells and can further strengthen their immunosuppressive potential (49, 85, 88). Also, vitamin D participates in redox balance by an activation in vitamin D receptor (VDR) and upregulation of antioxidant enzymes which can further preserve mitochondrial activity, decrease ROS accumulation, and support MSCs from oxidative injury (89, 90). Li *et al.* in their *in vivo* study found that VDR can protect against radiation-induced intestinal injury in mice via inhibition of intestinal crypt stem/progenitor cell apoptosis (91). It can improve autophagy and promote resistance to oxidative stress, specifically when combined with other metabolic modulators (59, 92-95). Vitamin D together with BM-MSCs has induced osteogenesis via enhancing expression of osteocalcin (OCN), ALP and Runx2 through VDR and a synergism with transforming growth factor-beta1 (TGF- β 1), bone morphogenetic protein 2 (BMP-2), dexamethasone and metformin to activate Wnt signaling pathway-catenin (Wnt/ β -catenin) and BMP2 signaling and a reduction in ROS (43, 44, 90, 96-99). Addition of vitamin D to human BM-MSCs has resulted in adipogenesis too due to a reduction in lipid accumulation and downregulation of adipocyte-specific genes and inhibition of transcription factor peroxisome proliferator-activated receptor gamma 2 (PPAR γ 2) (44, 100-102). This vitamin *in vitro* together with human periodontal ligament MSCs (PDL-MSCs) and with mouse BM-MSCs *in vivo* participated in bone fracture healing. Also, it showed strong anti-inflammatory effects *in vivo* in mouse model of acute and chronic diseases when used with lipopolysaccharide (LPS) and high fat diet. These changes happened via a decline in secretion of TNF- α , IL-1 β and IL-6 by blocking NF- κ B signaling; enhancing MSC-mediated suppression of CD4⁺ T cell proliferation; suppression of M1 macrophage differentiation; promotion in M2 macrophage differentiation and inhibition of M1 macrophage-mediated MSC migration in a dose-dependent manner (84, 85, 87). Vitamin D in *in vitro* addition to human and mouse BM-MSCs and *in vivo* with mouse model of high fat diet (HFD)-

induced osteoporosis could maintain redox balance based on activating VDR to upregulate superoxide dismutase 2 (SOD2) and preserving mitochondrial function (89, 90).

Recent studies illustrated the effect of the active form of vitamin D (vitamin D3) on maintenance of I-MSCs and inflammation in crypts. Supplementation of vitamin D3 and calcium to the diet can abrogate the adverse impact of NWD1 on I-MSC function. Vitamin D3 was shown to enhance the differentiation of I-MSCs into various specialized epithelial cells like Paneth, enterocytes, enteroendocrine and goblet cells. The crucial role of vitamin D in I-MSC regulation in Vdr knockout mouse model has been validated. Vitamin D3/VDR signalling pathway can protect I-MSCs and progenitor cells against DNA damage and apoptosis due to ionising radiation (50, 91, 103, 104).

Vitamin E

Vitamin E is fat-soluble antioxidant vitamin with eight known natural isoforms of four tocopherols and four tocotrienols, each designated as α , β , γ and δ based on the position of methyl groups on the chromanol ring. It is found in almost all the tissues in the human body, with the highest level in the adipose tissue and adrenal gland that plays a role as chain-breaking antioxidant, neutralizing free radicals and terminating chain reactions in the oxidation of polyunsaturated fatty acids. Vitamin E also can modulate cellular signal transduction through kinases, phosphatases, lipid mediators and transcription factors. Vitamin E was frequently utilized in primary cell culture to inhibit cell death and preserve cell activity when exposed to stress conditions. So commercial cell culture supplements that contain vitamin E are available (41).

Vitamin is essential for the maintenance of healthy skin. It is a group of eight compounds related in molecular structure enrolling four tocopherols and four tocotrienols (105). Pretreatment of MSCs with vitamin E was demonstrated to protect the stem cells from oxidative stress-induced damage and enhance the survival and preserve the regenerative capacity of MSCs (70). These findings can be promising in cartilage regeneration, as vitamin E-treated MSCs can promote chondrogenic differentiation and can slow the progression of osteoarthritis by an inhibition in apoptosis and senescence processes (106, 107). In *in vitro* preconditioning of MSCs and *in vivo* osteoarthritis rat models; an enhancement in MSC survival and cartilage regeneration was noticed. Preconditioning protected MSCs from oxidative stress (H₂O₂); downregulated apoptosis genes, reduced vascular endothelial growth factor

(VEGF) and lactate dehydrogenase (LDH) release, upregulated TGF- β , and increased proteoglycan content in cartilage (70, 106, 107). Vitamin E as a major fat-soluble antioxidant is vital to support cell membranes from lipid peroxidation and as a result can protect the MSC lipid bilayer from oxidative stress and has an anti-aging protection role and maintains MSC viability and activity, especially under the challenging conditions in tissue injuries (70, 108, 109). When vitamin E is added to human dental pulp MSCs (DP-MSCs), osteogenesis was noticed via enhancing osteogenic commitment (7, 98). Vitamin E when added *in vitro* to human DP-MSCs; porcine Ad-MSCs and rat BM-MSCs and *in vivo* in rat model of osteoarthritis (OA); it could protect from oxidative stress by preventing lipid peroxidation of cell membranes; reduction of oxidative stress and aging caused because of H₂O₂; maintenance of cell membrane integrity; protection against apoptosis and promoting survival and enhancing proliferation (70, 71, 110).

Vitamin K₂

Vitamin K₂ is found in animal foods such as liver, egg yolks, high fat dairy products, organ meats and fermented foods like sauerkraut and natto. The gut bacteria can also produce some K₂. It can help to decline the risk of cardiovascular diseases (CVDs) by preventing calcium buildup in arteries. It is divided into different subtypes like MK-4 and MK-7 (111). Vitamin K₂ may also support dental health by activating osteocalcin, a protein that helps stimulate new bone and dentin growth. Talk with a healthcare professional before adding vitamin K₂ supplements to the diet, especially if someone takes blood thinning medications like warfarin. Vitamin K₂ in combination with diet-induced obesity in mouse model and synergistically with vitamin D3 can lead to osteogenesis via enhancing the expression of osterix, osteocalcin and Runx2 (112).

Folic acid

Folic acid is a water-soluble B vitamin which has critical roles in neural tube development and neonatal gut maturation and overall gut health. Zheng *et al.* illustrated that folic acid promoted renewal, proliferation, differentiation of intestinal stem cells (ISCs) and demonstrated that folic acid could enhance ISC-driven epithelial regeneration in a β -adrenergic signaling-dependent manner and epithelial barrier integrity and homeostasis and could modulate epithelial function through the upregulation expression of tight junction components and nutrient transporters (113). Zhang *et al.* revealed that supplementation of maternal folic

acid can improve the intestinal health of porcine newborns through promoting the proliferation and differentiation of I-MSCs, increase in Ki67-positive cells and the expression of proliferation-related marker genes [(C-Myc, CyclinD1, and proliferating cell nuclear antigen (PCNA))] in piglets' intestinal stem cells (114).

Minerals

Essential minerals were exhibited to be integral to MSC-mediated repair. The differentiation of MSCs is prominently modulated by many minerals; thereby they have an important impact on tissue homeostasis and in regenerative medicine (3). Minerals such as iron, magnesium, zinc, calcium, selenium, etc. were highlighted as key micronutrients for stem cell activity (41, 46).

Iron

Iron is a key element in the metabolism of all living organisms and is found in two biologically oxidation states of the ferrous form (Fe²⁺) and the ferric form (Fe³⁺). It naturally exists in several foods, is added to some food products, and is available as a dietary supplement. Iron is an essential component in hemoglobin, an erythrocyte (red blood cell) protein that transfers oxygen from the lungs to the tissues. As a component of myoglobin as another protein can provide oxygen and support muscle metabolism. Iron is also essential for cellular functioning, neurological development, physical growth and hormonal synthesis (115-118).

Iron as an essential cofactor was demonstrated to have a prominent role for many proteins involved in critical biological processes, such as mitochondrial electron transport, DNA synthesis, and oxidative phosphorylation that make iron a fundamental metabolic currency. Its deficiency can impair intestinal mucosal structure and activity and affect intestinal stem cells and their differentiation potential (119). The dysregulated iron dynamics can lead to a significant decline in numbers and proliferative activity of intestinal stem cells as documented by a reduction in expression of the canonical stem cell marker leucine-rich repeat-containing G-protein coupled receptor 5 (Lgr5) and antigen Kiel 67 (Ki67)-positive proliferating cells. The correlation between nutrient deficiency and impaired proliferation of intestinal stem cells has been noticed in many models of malnutrition, revealing a potential common pathway in nutrient stress responses (120, 121).

It was suggested that iron delivered via iron oxide nanoparticles (IONPs) can enhance the migration of MSCs to the site of tissue injury and to facilitate

angiogenesis and wound healing; while IONP-labeled MSCs were also successfully employed to improve tissue repair in *in vivo* studies. These properties were successfully documented by magnetic resonance imaging (MRI) tracking (37, 40, 122-127) MSCs treated with iron and delivered to cells as IONPs displayed enhanced anti-inflammatory properties with a change in their cytokine profile toward the declined secretion of proinflammatory factors and a rise in anti-inflammatory cytokines (128).

Mouse model of dermal wounds; mouse model of hind limb ischemia for angiogenesis by using human Ad-MSCs and rat model of liver fibrosis utilizing rat BM-MSCs denoted to an improvement in wound healing and a reduction in liver fibrosis. It could enhance migration of MSCs to the site of tissue injury and led to an improvement in angiogenesis (122-124). Iron can illustrate a dualistic role too, as a decrease in its level can stimulate adaptive healing mechanisms, while excessive iron level can result in accumulation of toxic ROS that can underscore the importance of tightly regulated iron homeostasis (122, 129-131). Iron when treated with Ad-MSCs in mouse model of skin wounds and in mouse model of hindlimb ischemia (angiogenesis) and with BM-MSCs in rat model of liver fibrosis could improve wound healing and decrease liver fibrosis by enhancing migration of MSCs to the site of injury and lead to an improvement in angiogenesis (123-124).

Iron *in vitro* exposed to rat BM-MSCs and *in vivo* in rat model of laser burn wound could enhance anti-inflammatory properties through modification in cytokine profile and a decrease in pro-inflammatory factors (IL-2 and TNF- α) and a rise in secretion of anti-inflammatory cytokines of IL-4 and IL-10 (128). Iron in combination with human Ad-MSCs, human BM-MSCs and rat BM-MSCs revealed dualistic adaptive vs. toxic effects; while in low concentrations was associated with mild ROS for adaptive response [hypoxia-inducible factor 1 subunit alpha (HIF-1 α) stimulation] and in high concentrations was with toxic ROS (122, 129-131).

Magnesium

Magnesium is an abundant mineral in the body and a cofactor found in more than 300 enzyme systems that regulate various biochemical reactions in the body, like muscle and nerve function, protein synthesis, blood pressure regulation and blood glucose control. It is naturally found in various foods, available as a dietary supplement, added to food products, and seen in some medicines (e.g., antacids, laxatives) essential in oxidative phosphorylation, energy production and glycolysis.

It contributes to the structural development of bone and is necessary in RNA and DNA synthesis and the antioxidant glutathione. Magnesium also plays a role in the active transport of calcium and potassium ions across cell membranes to facilitate muscle contraction, nerve impulse conduction and normal heart rhythm (132).

Magnesium in addition to bone homeostasis is pivotal for both cartilage regeneration. Shimaya *et al.* reported that it can enhance the adhesion of synovial MSCs to injured sites, promote cartilage matrix synthesis, and induces apatite crystal growth on tissue surfaces as a key step in bone formation and remodeling (133, 134). *In vivo* study using a rabbit model of osteochondral defect with rabbit synovial MSCs; in *ex vivo* study employing osteochondral tissue and human synovial MSCs and in *in vitro* study of chondrogenesis assessment and cell adhesion utilizing human synovial MSCs and human Ad-MSCs, bone homeostasis and cartilage regeneration were supported. Magnesium could enhance MSC adhesion to defects; improve cartilage matrix synthesis and induced apatite crystal growth on scaffolds for mineralization (133, 134).

Magnesium has anti-inflammatory properties too that can significantly decrease secretion of proinflammatory cytokines from MSCs. It can simultaneously increase anti-inflammatory mediators such as prostaglandin E2 (PGE2) and IL-10 and modulate key inflammation-related transcription factors like NF- κ B to enhance the overall immunosuppressive activity of MSCs (135). Magnesium in combination with BM-MSCs could trigger osteogenesis by enhancing early differentiation, Notch and Wnt signaling and autophagy; while high concentrations were shown to impair late-stage mineralization (133, 136-141). When added to synovial MSCs, chondrogenesis was noted due to the support of differentiation and synthesis of cartilage-specific ECM via integrin-mediated signaling (133).

This mineral with synovial MSCs *in vivo* in rabbit model of osteochondral defect, in *ex vivo* with human synovial MSCs in osteochondral tissue; in *in vitro* with synovial MSCs in cell adhesion and chondrogenesis assessment; and *in vitro* with Ad-MSCs could support bone homeostasis and cartilage regeneration by promoting cartilage matrix synthesis; enhance MSC adhesion to defects and induce apatite crystal growth on scaffolds for mineralization (133, 134). Magnesium with murine MSCs and macrophages *in vitro* displayed strong anti-inflammatory and immunosuppressive impacts via a decrease in IL-1 β , IL-6; an increase in IL-10 and PGE2 and the modulation of nuclear factor NF- κ B

and signal transducer and activator of transcription-3 (STAT-3) (135).

Zinc

Zinc is an essential mineral that helps cell metabolism, growth, signaling and division. It is mainly stored in muscles and bones and is naturally present in some foods and is available as a dietary supplement. Zinc is involved in many aspects of cellular. It is required for the catalytic activity of hundreds of enzymes and is essential for protein and DNA synthesis, to enhance immune function and improve wound healing. Its homeostasis is maintained via absorption of zinc from the diet, excretion into the gastrointestinal tract, and reabsorption in gastrointestinal lumen. Zinc Supplementation can be beneficial in glycemic control, insulin resistance, inflammation and oxidative stress in diabetes too (142).

Zinc was described to improve the therapeutic efficiency of MSCs via enhancing the migration and adhesion of stem cells, which are important processes in homing of MSCs to the site of injury (4). Zinc-containing biomaterials were also tissue engineered to simultaneously improve angiogenesis and osteogenesis to support the complex tissue repair (143). *In vitro* scratch assays of human UC-MSCs and human Ad-MSCs with zinc promoted tissue repair via enhancing cell homing. It significantly improved migration and adhesion of MSCs and when scaffolds co-doped with copper and zinc, they promoted both angiogenesis and osteogenesis (4, 143).

Zinc is also essential in immune response regulation and its deficiency can be associated with an enhanced inflammation. Conversely, its supplementation has anti-inflammatory effects and can modulate the expression of genes involved in inflammatory signaling pathways of MSCs (144). It can enhance cellular defense via modulation of pathways that can promote the transcription of antioxidant genes (144, 145). Zinc when comes with Ad-MSCs can provide osteogenesis via enhancing matrix formation, osteoblast differentiation, and mineralization (143). Zinc and human UC-MSCs showed anti-inflammatory effects through upregulation of genes for cytokine receptor interactions and IL-17/TNF pathways. The deficiency was found to increase the TNF- α , IL-1 β and IL-8 (144).

Zinc in combination with human UC-MSCs could enhance antioxidant defense by modulating nuclear factor erythroid 2-related factor 2/sirtuin 3 (Nrf2/Sirt3) pathway to promote antioxidant gene transcription (144, 145). Zinc treated with rat Ad-MSCs could regulate histone deacetylation and telomeric stability by activating SIRT3 including histone deacetylase; functioning as a structural

component for zinc finger transcription factors and chromatin remodelers and an increase in telomerase reverse transcriptase (TERT) expression (146).

Calcium

Calcium is the most abundant metal and the 5th-most abundant element in the human body. Calcium ion (Ca²⁺) as an electrolyte plays a crucial role in the physiological and biochemical processes of cells: in signal transduction pathways where they play a role as a second messenger; in contraction of all muscle cell types; in neurotransmitter release from neurons; as cofactors in several enzymes and in fertilization. Its ions outside cells are important to maintain the potential difference across excitable cell membranes, protein synthesis, and bone formation (147). Presence of calcium with BM-MSCs and MSC immunoselected with stromal elements1 (STRO1) antibody provided the chance for osteogenesis. It happened because of upregulation of bone sialoprotein (BSP), OCN and ALP expressions and mediation through L-type calcium channels and calcium/calmodulin-dependent protein kinase II (CaMKII) signaling (148-150).

Selenium

Selenium is an ingredient in various multivitamins and dietary supplements, and in infant formulas. It is a component of the antioxidant enzymes glutathione peroxidase and thioredoxin reductase and in three deiodinase enzymes. Although trace amounts of selenium are essential in cellular function; but they can be toxic in even small doses and cause selenosis with symptoms of fatigue, diarrhea, hair loss, nail brittleness, joint pain or discoloration, headache, nausea, vomiting, tingling and fever (151). Selenium as a trace element is essential for antioxidant defense and specifically in nanoparticle forms that can present a strong, dose-dependent supportive influence against oxidative stress. At optimal levels, it can decrease intracellular ROS and activate signaling pathways to increase the expression of antioxidant enzymes and consequently improves MSCs viability (43).

Human embryonic stem cells-derived MSCs with selenium induced adipogenesis and osteogenesis. They happened through a change in MSC fate towards osteogenic lineage and an increase in ALP function and osteogenic gene transcription (43). Also it occurred via the suppression in adipocyte differentiation (43). Selenium addition to human ESC-derived MSCs and rat BM-MSCs showed a protecting effect against oxidative stress by activating c-Jun N-terminal kinase forkhead box O3a (JNK/FOXO3a) pathway and increasing

SOD and catalase (43, 152, 153). Selenium in combination with human BM-MSCs could support methylation cycle and chromatin integrity because as a component of selenoproteins, it can ensure the availability of S-adenosylmethionine (SAM), the universal methyl donor for DNA methyltransferases, histone H3K36me2/3 (DNMTs) and histone methyltransferases (HMTs) and decrease DNA damage indicators (micronuclei) by up to 58% (154).

Lithium chloride

Lithium is a fine, white, deliquescent granular or crystalline powder with a feeble alkaline taste. Its salts can influence the central nervous system (CNS) in a variety of ways and used to treat bipolar disorder (BPD) (14). Lithium chloride (LiCl) *in vitro* could enhance the proliferation and growth kinetic of Ad-MSCs and protect the cells from apoptosis via an increase in expression of the Bcl-2 gene and a significant decrease in Bax gene (155). LiCl intake with MSCs and a ketogenic diet can be successful in treatment of BPD. This happens by an improvement in neurogenesis and an increase in expression of neurotrophic factors such as brain derived neurotrophic factor (BDNF), nerve growth factor (NGF) and insulin-like growth factor-1 (IGF-1) that enhance the survival and differentiation of neurons (156).

Conclusion

Even many studies regarding the impact of vitamins and mineral components on proliferation, differentiation, homeostasis and regenerative capacity of body mesenchymal stem cells are *in vitro* and require *in vivo* validation; but the positive physiological effects of vitamins and minerals through several mechanisms in the literature reflect their pivotal role in regenerative medicine. It seems that improving nutritional intake and biochemical status of key micronutrients (e.g. vitamins and minerals) can be supportive for the efficacy of MSCs in therapies and immune function and subsequently the recovery and clinical outcome of patients. A proper food intake can particularly target the enhancement of stem cell activity, epigenetic regulation, redox-inflammatory pathways and cellular metabolism. In this review, the mitochondrial metabolic checkpoint regulating stem cell maintenance was highlighted. It was discussed how diet affects the metabolism of stem cells through the nutrient sensing pathways, thereby affecting stem cell fate decision and tissue regeneration.

Acknowledgment

It is essential to acknowledge the collaboration of Stem Cell Technology Center and The Department

of Clinical Nutrition of Shiraz University of Medical Sciences.

Authors' Contribution

Conceptualization: DM and SJM; Investigation: SM, ASM, MH and MN; Data curation: SM and MN; Writing original draft preparation: DM and MN; Writing review and editing: DM and SM; Supervision: DM and SJM. All authors have read and agreed to the published version of the manuscript.

Funding

No fund was received for this review.

Conflict of Interest

The authors declare no conflicts of interest.

References

- 1 Puca F, Fedele M, Rasio D, et al. Role of Diet in Stem and Cancer Stem Cells. *Int J Mol Sci.* 2022;23:8108. DOI: 10.3390/ijms23158108. PMID: 35897685.
- 2 Li WH, Yuan XY, Wang Z, et al. Nutrient sensing in intestinal stem cell: Linking dietary nutrients to cellular metabolic regulation. *World J Stem Cells.* 2025;17:107770. DOI: 10.4252/wjsc.v17.i7.107770. PMID: 40740539.
- 3 Kot M, Bronowicka-Adamska P, Tyszkiewicz-Czochara M. Regulation of Human Stem Cells by Functional Food Components: How Vitamins, Minerals and Phytochemicals Influence Mesenchymal Stem Cells' Fate and Function. *Nutrients.* 2025;17:3548. DOI: 10.3390/nu17223548. PMID: 41305599.
- 4 Sahibdad I, Khalid S, Chaudhry GR, et al. Zinc enhances the cell adhesion, migration, and self-renewal potential of human umbilical cord derived mesenchymal stem cells. *World J Stem Cells.* 2023;15:751-767. DOI: 10.4252/wjsc.v15.i7.751. PMID: 37545753.
- 5 Escobar LM, Bendahan Z, Bayona A, et al. Effect of Vitamins D and E on the Proliferation, Viability, and Differentiation of Human Dental Pulp Stem Cells: An In Vitro Study. *Int J Dent.* 2020;2020:8860840. DOI: 10.1155/2020/8860840. PMID: 32676112.
- 6 Saud B, Malla R, Shrestha K. A Review on the Effect of Plant Extract on Mesenchymal Stem Cell Proliferation and Differentiation. *Stem Cells Int.* 2019;2019:7513404. DOI: 10.1155/2019/7513404. PMID: 31428160.
- 7 Van Pham P, Tran NY, Phan NL, et al. Vitamin C stimulates human gingival stem cell proliferation and expression of pluripotent markers. *In Vitro Cell Dev Biol Anim.* 2016;52:218-27. DOI:

- 10.1007/s11626-015-9963-2. PMID: 26487430.
- 8 D'Aniello C, Cermola F, Patriarca EJ, et al. Vitamin C in Stem Cell Biology: Impact on Extracellular Matrix Homeostasis and Epigenetics. *Stem Cells Int.* 2017;2017:8936156. DOI: 10.1155/2017/8936156. PMID: 28512473.
- 9 Khillan JS. Vitamin A/retinol and maintenance of pluripotency of stem cells. *Nutrients.* 2014;6:1209-22. DOI: 10.3390/nu6031209. PMID: 24662164.
- 10 Barthez M, Song Z, Wang CL, et al. Stem Cell Metabolism and Diet. *Curr Stem Cell Rep.* 2020;6:119-125. DOI: 10.1007/s40778-020-00180-4. PMID: 33777658.
- 11 Fekete N, Rojewski MT, Lotfi R, et al. Essential components for ex vivo proliferation of mesenchymal stromal cells. *Tissue Eng Part C Methods.* 2014;20:129-39. DOI: 10.1089/ten.TEC.2013.0061. PMID: 23713576.
- 12 Adefegha SA. Functional Foods and Nutraceuticals as Dietary Intervention in Chronic Diseases; Novel Perspectives for Health Promotion and Disease Prevention. *J Diet Suppl.* 2018;15:977-1009. DOI: 10.1080/19390211.2017.1401573. PMID: 29281341.
- 13 Mehrabani D, Masoumi SJ, Masoumi AS, et al. Role of Diet in Mesenchymal Stem Cells' Function: A Review. *Int J Nutr Sci.* 2023;8:9-19. DOI: 10.30476/IJNS.2023.97788.1221.
- 14 Homayoon M, Mehrabani D, Edalatmanesh MA, et al. In Vitro Effect of Lithium Chloride on Adipose Tissue Derived Stem Cells Proliferation and Growth Kinetic. *Iran J Med Sci.* 2026;51:445-454. DOI: 10.30476/ijms.2026.108735.4372. PMID: 42370021.
- 15 Ho J, Puoplo N, Pokharel N, et al. Nutrigenomic underpinnings of intestinal stem cells in inflammatory bowel disease and colorectal cancer development. *Front Genet.* 2024;15:1349717. DOI: 10.3389/fgene.2024.1349717. PMID: 39280096.
- 16 Yadav UP, Singh T, Kumar P, et al. Metabolic Adaptations in Cancer Stem Cells. *Front Oncol.* 2020;10:1010. DOI: 10.3389/fonc.2020.01010. PMID: 32670883.
- 17 Lu V, Roy IJ, Teitell MA. Nutrients in the fate of pluripotent stem cells. *Cell Metab.* 2021;33:2108-2121. DOI: 10.1016/j.cmet.2021.09.013. PMID: 34644538.
- 18 Baksh SC, Todorova PK, Gur-Cohen S, et al. Extracellular serine controls epidermal stem cell fate and tumour initiation. *Nat Cell Biol.* 2020;22:779-790. DOI: 10.1038/s41556-020-0525-9. PMID: 32451440.
- 19 Bifari F, Dolci S, Bottani E, et al. Complete neural stem cell (NSC) neuronal differentiation requires a branched chain amino acids-induced persistent metabolic shift towards energy metabolism. *Pharmacol Res.* 2020;158:104863. DOI: 10.1016/j.phrs.2020.104863. PMID: 32407957.
- 20 Panopoulos AD, Yanes O, Ruiz S, et al. The metabolome of induced pluripotent stem cells reveals metabolic changes occurring in somatic cell reprogramming. *Cell Res.* 2012;22:168-77. DOI: 10.1038/cr.2011.177. PMID: 22064701.
- 21 Yilmaz ÖH, Katajisto P, Lamming DW, et al. mTORC1 in the Paneth cell niche couples intestinal stem-cell function to calorie intake. *Nature.* 2012;486:490-5. DOI: 10.1038/nature11163. PMID: 22722868.
- 22 Smith J, Ladi E, Mayer-Proschel M, et al. Redox state is a central modulator of the balance between self-renewal and differentiation in a dividing glial precursor cell. *Proc Natl Acad Sci U S A.* 2000;97:10032-7. DOI: 10.1073/pnas.170209797. PMID: 10944195.
- 23 Ghobadi F, Mehrabani D, Mehrabani G. Regenerative potential of endometrial stem cells: a mini review. *World J Plast Surg.* 2015;4:3-8. PMID: 25606470.
- 24 Mehrabani D, Sholehvar F, Yaghmaei P, et al. The impact of acemannan, an extracted product from Aloe vera, on proliferation of dental pulp stem cells and healing of mandibular defects in rabbits. *Am J Stem Cells.* 2024;13:75-86. DOI: 10.62347/UAFC3719. PMID: 38765804.
- 25 Mehrabani D, Rahmanifar F, Mellinejad M, et al. Isolation, culture, characterization, and adipogenic differentiation of heifer endometrial mesenchymal stem cells. *Comp Clin Pathol.* 2015;24:1159-64. DOI: 10.1007/s00580-014-2053-0.
- 26 Hosseini-Asl SK, Mehrabani D, Karimi-Busheri F. Therapeutic Effect of Mesenchymal Stem Cells in Ulcerative Colitis: A Review on Achievements and Challenges. *J Clin Med.* 2020;9:3922. DOI: 10.3390/jcm9123922. PMID: 33287220.
- 27 Irani D, Mehrabani D, Karimi-Busheri F. Mesenchymal Stem Cells in Regenerative Medicine, Possible Applications in The Restoration of Spermatogenesis: A Review. *Cell J.* 2024;26:169-184. DOI: 10.22074/cellj.2024.2015141.1442. PMID: 38628090.
- 28 Kaboodkhani R, Mehrabani D, Karimi-Busheri F. Achievements and Challenges in Transplantation of Mesenchymal Stem Cells in Otorhinolaryngology. *J Clin Med.* 2021;10:2940. DOI: 10.3390/jcm10132940. PMID: 34209041.
- 29 Hajihoseini M, Vahdati A, Ebrahim Hosseini S, et al. Induction of spermatogenesis after stem cell

- therapy of azoospermic guinea pigs. *Vet Arhiv*. 2017;87:333-50. DOI: 10.24099/vet.arhiv.151209.
- 30 Mehrabani D, Khodakaram-Tafti A, Shaterzadeh-Yazdi H, et al. Comparison of the regenerative effect of adipose-derived stem cells, fibrin glue scaffold, and autologous bone graft in experimental mandibular defect in rabbit. *Dent Traumatol*. 2018;34:413-420. DOI: 10.1111/edt.12435. PMID: 30187637.
 - 31 Mehrabani D, Mahdiyar P, Torabi K, et al. Growth kinetics and characterization of human dental pulp stem cells: Comparison between third molar and first premolar teeth. *J Clin Exp Dent*. 2017;9:e172-e177. DOI: 10.4317/jced.52824. PMID: 28210430.
 - 32 Nazempour M, Mehrabani D, Mehdinavaz-Aghdam R, et al. The effect of allogenic human Wharton's jelly stem cells seeded onto acellular dermal matrix in healing of rat burn wounds. *J Cosmet Dermatol*. 2020;19:995-1001. DOI: 10.1111/jocd.13109. PMID: 31556227.
 - 33 Ghobadi F, Rahmanifar F, Mehrabani D, et al. Endometrial mesenchymal stem stromal cells in mature and immature sheep: An in vitro study. *Int J Reprod Biomed*. 2018;16:83-92. PMID: 29675492.
 - 34 Mehrabani D, Nazarabadi RB, Kasraeian M, et al. Growth Kinetics, Characterization, and Plasticity of Human Menstrual Blood Stem Cells. *Iran J Med Sci*. 2016;41:132-9. PMID: 26989284.
 - 35 Karimaghai N, Tamadon A, Rahmanifar F, et al. Spermatogenesis after transplantation of adipose tissue-derived mesenchymal stem cells in busulfan-induced azoospermic hamster. *Iran J Basic Med Sci*. 2018;21:660-667. DOI: 10.22038/IJBMS.2018.29040.7010. PMID: 30140403.
 - 36 Mehrabani D, Khodabandeh Z. Effect of bone marrow-derived mesenchymal stem cells on changes of serum levels of $\text{tnf-}\alpha$ and locomotor function after spinal cord injury in mice. *J Med Sci*. 2016;16:16-24. DOI: 10.3923/jms.2016.16.24.
 - 37 Hashemi SS, Niknam Z, Mojtaba Zebarjad S, et al. Tracking the healing effect of human Wharton's jelly stem cells labeled with superparamagnetic iron oxide nanoparticles seeded onto polyvinyl alcohol/chitosan/carbon nanotubes in burn wounds by MRI and Prussian blue staining. *Biomed Mater*. 2025;20. DOI: 10.1088/1748-605X/ad9fc6. PMID: 39681069.
 - 38 Mehrabani D, Babazadeh M, Tanideh N, et al. The Healing Effect of Adipose-Derived Mesenchymal Stem Cells in Full-thickness Femoral Articular Cartilage Defects of Rabbit. *Int J Organ Transplant Med*. 2015;6:165-75. PMID: 26576262.
 - 39 Mehrabani D, Khajehahmadi Z, Tajik P, et al. Regenerative Effect of Bone Marrow-derived Mesenchymal Stem Cells in Thioacetamide-induced Liver Fibrosis of Rats. *Arch Razi Inst*. 2019;74:279-286. DOI: 10.22092/ari.2018.110029.1120. PMID: 31592593.
 - 40 Zare S, Mehrabani D, Jalli R, et al. MRI-Tracking of Dental Pulp Stem Cells In Vitro and In Vivo Using Dextran-Coated Superparamagnetic Iron Oxide Nanoparticles. *J Clin Med*. 2019;8:1418. DOI: 10.3390/jcm8091418. PMID: 31505807.
 - 41 Godoy-Parejo C, Deng C, Zhang Y, et al. Roles of vitamins in stem cells. *Cell Mol Life Sci*. 2020;77:1771-1791. DOI: 10.1007/s00018-019-03352-6. PMID: 31676963.
 - 42 Wang SH, Lee SP, Yang CW, et al. Surface modification of biodegradable Mg-based scaffolds for human mesenchymal stem cell proliferation and osteogenic differentiation. *Materials*. 2021;14:441. DOI: 10.3390/ma14020441.
 - 43 Fatima S, Alfrayh R, Alrashed M, et al. Selenium Nanoparticles by Moderating Oxidative Stress Promote Differentiation of Mesenchymal Stem Cells to Osteoblasts. *Int J Nanomedicine*. 2021;16:331-343. DOI: 10.2147/IJN.S285233. PMID: 33488075.
 - 44 Borojević A, Jauković A, Kukolj T, et al. Vitamin D3 Stimulates Proliferation Capacity, Expression of Pluripotency Markers, and Osteogenesis of Human Bone Marrow Mesenchymal Stromal/Stem Cells, Partly through SIRT1 Signaling. *Biomolecules*. 2022;12:323. DOI: 10.3390/biom12020323. PMID: 35204824.
 - 45 Sharifi S, Moghaddam FA, Abedi A, et al. Phytochemicals impact on osteogenic differentiation of mesenchymal stem cells. *BioFactors*. 2020;46:874-93. DOI: 10.1002/biof.1682.
 - 46 Zong Q, Bundkirchen K, Neunaber C, et al. Are the Properties of Bone Marrow-Derived Mesenchymal Stem Cells Influenced by Overweight and Obesity? *Int J Mol Sci*. 2023;24:4831. DOI: 10.3390/ijms24054831. PMID: 36902259.
 - 47 Özcan SG, Stanko C, Szymanski L, et al. Revisiting retinoic acid in AML therapy: mechanisms of action and rational combination strategies. *Br J Cancer*. 2026. DOI: 10.1038/s41416-026-03512-z. PMID: 42393459.
 - 48 de Witte SFH, Merino AM, Franquesa M, et al. Cytokine treatment optimises the immunotherapeutic effects of umbilical cord-derived MSC for treatment of inflammatory liver disease. *Stem Cell Res Ther*. 2017;8:140. DOI: 10.1186/s13287-017-0590-6. PMID: 28595619.

- 49 Di Tinco R, Bertani G, Pisciotto A, et al. Role of PD-L1 in licensing immunoregulatory function of dental pulp mesenchymal stem cells. *Stem Cell Res Ther.* 2021;12:598. DOI: 10.1186/s13287-021-02664-4. PMID: 34863286.
- 50 Shi R, Wang B. Nutrient metabolism in regulating intestinal stem cell homeostasis. *Cell Prolif.* 2024;57:e13602. DOI: 10.1111/cpr.13602. PMID: 38386338.
- 51 Wang Z, Li J, Wang Y, et al. Dietary vitamin A affects growth performance, intestinal development, and functions in weaned piglets by affecting intestinal stem cells. *J Anim Sci.* 2020;98:skaa020. DOI: 10.1093/jas/skaa020. PMID: 31955210.
- 52 Ghosh SK, Bandyopadhyay D, Panja S. B vitamins in dermatology. *Clin Dermatol.* 2026;44:223-240. DOI: 10.1016/j.clindermatol.2026.02.006. PMID: 41692080.
- 53 Ok JS, Song SB, Hwang ES. Enhancement of Replication and Differentiation Potential of Human Bone Marrow Stem Cells by Nicotinamide Treatment. *Int J Stem Cells.* 2018;11:13-25. DOI: 10.15283/ijsc18033. PMID: 29699388.
- 54 Ancel S, Michaud J, Migliavacca E, et al. Nicotinamide and pyridoxine stimulate muscle stem cell expansion and enhance regenerative capacity during aging. *J Clin Invest.* 2024;134:e163648. DOI: 10.1172/JCI163648. PMID: 39531334.
- 55 Kato N, Yang Y, Khedara A, et al. Potential Role of Vitamin B6 as an Antioxidant via Pyridoxal-5'-Phosphate-Dependent Metabolic Pathways and Subsequent Activation of Nrf2 Signaling. *Nutrients.* 2026;18:1499. DOI: 10.3390/nu18101499. PMID: 42196957.
- 56 Shotorbani BB, André H, Barzegar A, et al. Cell sheet biofabrication by co-administration of mesenchymal stem cells secretome and vitamin C on thermoresponsive polymer. *J Mater Sci Mater Med.* 2018;29:170. DOI: 10.1007/s10856-018-6180-z. PMID: 30392027.
- 57 Sarkar N, Morton H, Bose S. Effects of vitamin C on osteoblast proliferation and osteosarcoma inhibition using plasma coated hydroxyapatite on titanium implants. *Surf Coat Technol.* 2020;394:125793. DOI: 10.1016/j.surfcoat.2020.125793. PMID: 32612317.
- 58 Qu YN, Zhang L, Wang T, et al. Vitamin C Treatment Rescues Prelamin A-Induced Premature Senescence of Subchondral Bone Mesenchymal Stem Cells. *Stem Cells Int.* 2020;2020:3150716. DOI: 10.1155/2020/3150716. PMID: 32322277.
- 59 Yang M, Teng S, Ma C, et al. Ascorbic acid inhibits senescence in mesenchymal stem cells through ROS and AKT/mTOR signaling. *Cytotechnology.* 2018;70:1301-1313. DOI: 10.1007/s10616-018-0220-x. PMID: 29777434.
- 60 Lee ES, Chan J, Shuter B, et al. Microgel iron oxide nanoparticles for tracking human fetal mesenchymal stem cells through magnetic resonance imaging. *Stem Cells.* 2009;27:1921-31. DOI: 10.1002/stem.112. PMID: 19544438.
- 61 Zhao L, Tang M, Weir MD, et al. Osteogenic media and rhBMP-2-induced differentiation of umbilical cord mesenchymal stem cells encapsulated in alginate microbeads and integrated in an injectable calcium phosphate-chitosan fibrous scaffold. *Tissue Eng Part A.* 2011;17:969-79. DOI: 10.1089/ten.TEA.2010.0521. PMID: 21091340.
- 62 Castano-Izquierdo H, Alvarez-Barreto J, van den Dolder J, et al. Pre-culture period of mesenchymal stem cells in osteogenic media influences their in vivo bone forming potential. *J Biomed Mater Res A.* 2007;82:129-38. DOI: 10.1002/jbm.a.31082. PMID: 17269144.
- 63 Cimmino L, Neel BG, Aifantis I. Vitamin C in Stem Cell Reprogramming and Cancer. *Trends Cell Biol.* 2018;28:698-708. DOI: 10.1016/j.tcb.2018.04.001. PMID: 29724526.
- 64 Wei C, Liu X, Tao J, et al. Effects of vitamin C on characteristics retaining of in vitro-cultured mouse adipose-derived stem cells. *In Vitro Cell Dev Biol Anim.* 2014;50:75-86. DOI: 10.1007/s11626-013-9673-6. PMID: 23949782.
- 65 Serpooshan V, Julien M, Nguyen O, et al. Reduced hydraulic permeability of three-dimensional collagen scaffolds attenuates gel contraction and promotes the growth and differentiation of mesenchymal stem cells. *Acta Biomater.* 2010;6:3978-87. DOI: 10.1016/j.actbio.2010.04.028. PMID: 20451675.
- 66 Virdee SS, Bashir N, Camilleri J, et al. Exploiting Dentine Matrix Proteins in Cell-Free Approaches for Periradicular Tissue Engineering. *Tissue Eng Part B Rev.* 2022;28:707-732. DOI: 10.1089/ten.TEB.2021.0074. PMID: 34309453.
- 67 Carr AC, Maggini S. Vitamin C and Immune Function. *Nutrients.* 2017;9:1211. DOI: 10.3390/nu9111211. PMID: 29099763.
- 68 Lee Chong T, Ahearn EL, Cimmino L. Reprogramming the Epigenome With Vitamin C. *Front Cell Dev Biol.* 2019;7:128. DOI: 10.3389/fcell.2019.00128. PMID: 31380368.
- 69 Yang M, Teng S, Ma C, et al. Ascorbic acid inhibits senescence in mesenchymal stem cells through ROS and AKT/mTOR signaling. *Cytotechnology.* 2018;70:1301-1313. DOI: 10.1007/s10616-018-0220-x. PMID: 29777434.

- 10.1007/s10616-018-0220-x. PMID: 29777434.
- 70 Bhatti FU, Mehmood A, Latief N, et al. Vitamin E protects rat mesenchymal stem cells against hydrogen peroxide-induced oxidative stress in vitro and improves their therapeutic potential in surgically-induced rat model of osteoarthritis. *Osteoarthritis Cartilage*. 2017;25:321-331. DOI: 10.1016/j.joca.2016.09.014. PMID: 27693502.
 - 71 Bhatti FUR, Kim SJ, Yi AK, et al. Cytoprotective role of vitamin E in porcine adipose-tissue-derived mesenchymal stem cells against hydrogen-peroxide-induced oxidative stress. *Cell Tissue Res*. 2018;374:111-120. DOI: 10.1007/s00441-018-2857-3. PMID: 29951700.
 - 72 Wang Y, Singh A, Xu P, et al. Expansion and osteogenic differentiation of bone marrow-derived mesenchymal stem cells on a vitamin C functionalized polymer. *Biomaterials*. 2006;27:3265-73. DOI: 10.1016/j.biomaterials.2006.01.036. PMID: 16494940.
 - 73 Burgess KA, Herrick AL, Watson REB. Systemic sclerosis skin is a primed microenvironment for soft tissue calcification-a hypothesis. *Rheumatology (Oxford)*. 2021;60:2517-2527. DOI: 10.1093/rheumatology/keab156. PMID: 33585894.
 - 74 Rahman F, Bordignon B, Culerrier R, et al. Ascorbic acid drives the differentiation of mesoderm-derived embryonic stem cells. Involvement of p38 MAPK/CREB and SVCT2 transporter. *Mol Nutr Food Res*. 2017;61. DOI: 10.1002/mnfr.201600506. PMID: 27957816.
 - 75 Barlian A, Judawisastra H, Alfarafisa NM, et al. Chondrogenic differentiation of adipose-derived mesenchymal stem cells induced by L-ascorbic acid and platelet rich plasma on silk fibroin scaffold. *PeerJ*. 2018;6:e5809. DOI: 10.7717/peerj.5809. PMID: 30488014.
 - 76 Sakurai R, Singh H, Wang Y, et al. Effect of Perinatal Vitamin D Deficiency on Lung Mesenchymal Stem Cell Differentiation and Injury Repair Potential. *Am J Respir Cell Mol Biol*. 2021;65:521-531. DOI: 10.1165/rcmb.2020-0183OC. PMID: 34126864.
 - 77 Rathinavelu S, Guidry-Elizondo C, Banu J. Molecular Modulation of Osteoblasts and Osteoclasts in Type 2 Diabetes. *J Diabetes Res*. 2018;2018:6354787. DOI: 10.1155/2018/6354787. PMID: 30525054.
 - 78 Schröder-Heurich B, von Hardenberg S, Brodowski L, et al. Vitamin D improves endothelial barrier integrity and counteracts inflammatory effects on endothelial progenitor cells. *FASEB J*. 2019;33:9142-9153. DOI: 10.1096/fj.201802750RR. PMID: 31084577.
 - 79 Mortazavi SMJ, Kadivar F, Zare S, et al. The Role of Adipose Tissue-Derived Stem Cells together with Vitamin C on Survival of Rats with Acute Radiation Syndrome. *J Biomed Phys Eng*. 2026;16:87-96. DOI: 10.31661/jbpe.v0i0.2310-1680. PMID: 42057983.
 - 80 Bourebaba L, Michalak I, Röcken M, et al. Cladophora glomerata methanolic extract decreases oxidative stress and improves viability and mitochondrial potential in equine adipose derived mesenchymal stem cells (ASCs). *Biomed Pharmacother*. 2019;111:6-18. DOI: 10.1016/j.biopha.2018.12.020. PMID: 30553132.
 - 81 Thienpont B, Steinbacher J, Zhao H, et al. Tumour hypoxia causes DNA hypermethylation by reducing TET activity. *Nature*. 2016;537:63-68. DOI: 10.1038/nature19081. PMID: 27533040.
 - 82 Guo Y, Yan Y, Zhao L, et al. Epigenetic Mechanisms of Vitamin D in the Aging Process: A Narrative Review. *Int J Mol Sci*. 2026;27:6578. DOI: 10.3390/ijms27156578. PMID: 42589236.
 - 83 Li W, Zimmerman SE, Peregrina K, et al. The nutritional environment determines which and how intestinal stem cells contribute to homeostasis and tumorigenesis. *Carcinogenesis*. 2019;40:937-946. DOI: 10.1093/carcin/bgz106. PMID: 31169292.
 - 84 Wasnik S, Rundle CH, Baylink DJ, et al. 1,25-Dihydroxyvitamin D suppresses M1 macrophages and promotes M2 differentiation at bone injury sites. *JCI Insight*. 2018;3:e98773. DOI: 10.1172/jci.insight.98773. PMID: 30185660.
 - 85 Behm C, Blufstein A, Gahn J, et al. 1,25(OH)2D3 Differently Affects Immunomodulatory Activities of Mesenchymal Stem Cells Depending on the Presence of TNF- α , IL-1 β and IFN- γ . *J Clin Med*. 2019;8:2211. DOI: 10.3390/jcm8122211. PMID: 31847340.
 - 86 Marcotorchino J, Gouranton E, Romier B, et al. Vitamin D reduces the inflammatory response and restores glucose uptake in adipocytes. *Mol Nutr Food Res*. 2012;56:1771-82. DOI: 10.1002/mnfr.201200383. PMID: 23065818.
 - 87 Karkeni E, Marcotorchino J, Tourniaire F, et al. Vitamin D limits chemokine expression in adipocytes and macrophage migration in vitro and in male mice. *Endocrinology*. 2015;156:1782-93. DOI: 10.1210/en.2014-1647. PMID: 25730105.
 - 88 Ahmadi P, Yan M, Bauche A, et al. Human dental pulp cells modulate CD8+ T cell proliferation and efficiently degrade extracellular ATP to adenosine in vitro. *Cell Immunol*. 2022;380:104589. DOI: 10.1016/j.cellimm.2022.104589. PMID: 36084401.
 - 89 Klotz B, Mentrup B, Regensburger M, et al.

- 1,25-dihydroxyvitamin D3 treatment delays cellular aging in human mesenchymal stem cells while maintaining their multipotent capacity. *PLoS One*. 2012;7:e29959. DOI: 10.1371/journal.pone.0029959. PMID: 22242193.
- 90 Chen J, Kuang S, Cen J, et al. Multiomics profiling reveals VDR as a central regulator of mesenchymal stem cell senescence with a known association with osteoporosis after high-fat diet exposure. *Int J Oral Sci*. 2024;16:41. DOI: 10.1038/s41368-024-00309-9. PMID: 38777841.
 - 91 Li W, Lin Y, Luo Y, et al. Vitamin D Receptor Protects against Radiation-Induced Intestinal Injury in Mice via Inhibition of Intestinal Crypt Stem/Progenitor Cell Apoptosis. *Nutrients*. 2021;13:2910. DOI: 10.3390/nu13092910. PMID: 34578802.
 - 92 Khorraminejad-Shirazi M, Sani M, Talaei-Khozani T, et al. AICAR and nicotinamide treatment synergistically augment the proliferation and attenuate senescence-associated changes in mesenchymal stromal cells. *Stem Cell Res Ther*. 2020;11:45. DOI: 10.1186/s13287-020-1565-6. PMID: 32014016.
 - 93 Cruciani S, Garroni G, Pala R, et al. Metformin and Vitamin D Modulate Inflammation and Autophagy during Adipose-Derived Stem Cell Differentiation. *Int J Mol Sci*. 2021;22:6686. DOI: 10.3390/ijms22136686. PMID: 34206506.
 - 94 Cruciani S, Garroni G, Pala R, et al. Metformin and vitamin D modulate adipose-derived stem cell differentiation towards the beige phenotype. *Adipocyte*. 2022;11:356-365. DOI: 10.1080/21623945.2022.2085417. PMID: 35734882.
 - 95 Le Pelletier L, Mantecon M, Gorwood J, et al. Metformin alleviates stress-induced cellular senescence of aging human adipose stromal cells and the ensuing adipocyte dysfunction. *Elife*. 2021;10:e62635. DOI: 10.7554/eLife.62635. PMID: 34544550.
 - 96 Valenti MT, Dalle Carbonare L, Mottes M. Osteogenic Differentiation in Healthy and Pathological Conditions. *Int J Mol Sci*. 2016;18:41. DOI: 10.3390/ijms18010041. PMID: 28035992.
 - 97 Kim S, Lee S, Kim K. Bone Tissue Engineering Strategies in Co-Delivery of Bone Morphogenetic Protein-2 and Biochemical Signaling Factors. *Adv Exp Med Biol*. 2018;1078:233-244. DOI: 10.1007/978-981-13-0950-2_12. PMID: 30357626.
 - 98 Song I, Kim BS, Kim CS, et al. Effects of BMP-2 and vitamin D3 on the osteogenic differentiation of adipose stem cells. *Biochem Biophys Res Commun*. 2011;408:126-31. DOI: 10.1016/j.bbrc.2011.03.135. PMID: 21463608.
 - 99 Lu CL, Shyu JF, Wu CC, et al. Association of Anabolic Effect of Calcitriol with Osteoclast-Derived Wnt 10b Secretion. *Nutrients*. 2018;10:1164. DOI: 10.3390/nu10091164. PMID: 30149605.
 - 100 Duque G, Rivas D. Alendronate has an anabolic effect on bone through the differentiation of mesenchymal stem cells. *J Bone Miner Res*. 2007;22:1603-11. DOI: 10.1359/jbmr.070701. PMID: 17605634.
 - 101 Muruganandan S, Roman AA, Sinal CJ. Adipocyte differentiation of bone marrow-derived mesenchymal stem cells: cross talk with the osteoblastogenic program. *Cell Mol Life Sci*. 2009;66:236-53. DOI: 10.1007/s00018-008-8429-z. PMID: 18854943.
 - 102 Pesarini JR, Oliveira RJ, Pessatto LR, et al. Vitamin D: Correlation with biochemical and body composition changes in a southern Brazilian population and induction of cytotoxicity in mesenchymal stem cells derived from human adipose tissue. *Biomed Pharmacother*. 2017;91:861-871. DOI: 10.1016/j.biopha.2017.05.013. PMID: 28501775.
 - 103 Peregrina K, Houston M, Daroqui C, et al. Vitamin D is a determinant of mouse intestinal Lgr5 stem cell functions. *Carcinogenesis*. 2015;36:25-31. DOI: 10.1093/carcin/bgu221. PMID: 25344836.
 - 104 Sittipo P, Kim HK, Han J, et al. Vitamin D3 suppresses intestinal epithelial stemness via ER stress induction in intestinal organoids. *Stem Cell Res Ther*. 2021;12:285. DOI: 10.1186/s13287-021-02361-2. PMID: 33985576.
 - 105 Ali M, Sheikh Abdul Kadir SH, Ibrahim E, et al. Tocotrienol-rich fraction of vitamin E in diabetes and cardiovascular disease: From molecular mechanisms to clinical application. *Int J Cardiol Cardiovasc Risk Prev*. 2026;30:200686. DOI: 10.1016/j.ijcrp.2026.200686. PMID: 42565029.
 - 106 Hu C, Li L. Preconditioning influences mesenchymal stem cell properties in vitro and in vivo. *J Cell Mol Med*. 2018;22:1428-1442. DOI: 10.1111/jcmm.13492. PMID: 29392844.
 - 107 Kornicka K, Babiarczuk B, Krzak J, et al. The effect of a sol-gel derived silica coating doped with vitamin E on oxidative stress and senescence of human adipose-derived mesenchymal stem cells (AMSCs). *RSC Adv*. 2016;6:29524-37. DOI: 10.1039/c6ra00029k.
 - 108 Casado-Díaz A, Túnez-Fiñana I, Mata-Granados JM, et al. Serum from postmenopausal women treated with a by-product of olive-oil extraction process stimulates osteoblastogenesis and

- inhibits adipogenesis in human mesenchymal stem-cells (MSC). *Exp Gerontol.* 2017;90:71-78. DOI: 10.1016/j.exger.2017.01.024. PMID: 28167238.
- 109 Casati L, Pagani F, Limonta P, et al. Beneficial effects of δ -tocotrienol against oxidative stress in osteoblastic cells: studies on the mechanisms of action. *Eur J Nutr.* 2020;59(5):1975-1987. DOI: 10.1007/s00394-019-02047-9. PMID: 31280345.
 - 110 El Alami M, Viña-Almunia J, Gambini J, et al. Activation of p38, p21, and NRF-2 mediates decreased proliferation of human dental pulp stem cells cultured under 21% O₂. *Stem Cell Reports.* 2014;3:566-73. DOI: 10.1016/j.stemcr.2014.08.002. PMID: 25358785.
 - 111 Tuxen HN, Zwisler L, Linneberg A. The History of the Discovery of Vitamin K. *Basic Clin Pharmacol Toxicol.* 2026;139:e70283. DOI: 10.1111/bcpt.70283. PMID: 42575715.
 - 112 Poon CC, Li RW, Seto SW, et al. In vitro vitamin K(2) and 1 α ,25-dihydroxyvitamin D(3) combination enhances osteoblasts anabolism of diabetic mice. *Eur J Pharmacol.* 2015;767:30-40. DOI: 10.1016/j.ejphar.2015.09.048. PMID: 26452518.
 - 113 Zheng Y, Han Y, Zhang Y, et al. Folic acid enhances intestinal stem cell-mediated epithelial regeneration depending on β -adrenergic receptor signaling. *Food Funct.* 2026. DOI: 10.1039/d6fo01675h. Epub ahead of print. PMID: 42300427.
 - 114 Zhang Y, Zhang X, Chen J, et al. Maternal Folic Acid Supplementation Improves the Intestinal Health of Offspring Porcine by Promoting the Proliferation and Differentiation of Intestinal Stem Cells. *Animals (Basel).* 2023;13:3092. DOI: 10.3390/ani13193092. PMID: 37835698.
 - 115 Fujii T. Iron Deficiency in Athletes. *Biol Trace Elem Res.* 2026. DOI: 10.1007/s12011-026-05289-x. PMID: 42573718.
 - 116 Pasalar M, Mehrabani D, Afrasiabi A, Mehravar Z, et al. Prevalence of thalassaemia, iron-deficiency anaemia and glucose-6-phosphate dehydrogenase deficiency among Arab migrating nomad children, southern Islamic Republic of Iran. *East Mediterr Health J.* 2014;20:726-31. PMID: 25601811.
 - 117 Karimi M, Mehrabani D, Pasalar M, et al. Thalassemia, Iron And G6pd Deficiency In Lor Migrating Nomad Children, Southern Iran. *Iran Red Crescent Med J.* 2010;12:441-445.
 - 118 Mehrabani D, Karimi M, Pasalar M, et al. Frequency Of Thalassemia, Iron And Glucose-6phosphate Dehydrogenase Deficiency Among Turkish Migrating Nomad Children In Southern Iran. *Acta Med Iran.* 2008;47:1-5.
 - 119 Xu Y, Zhao J, Jiang S, et al. Iron Deficiency Inhibits the Proliferation of Intestinal Stem Cells and Induces Their Differentiation to Enterocytes. *Nutrients.* 2026;18:392. DOI: 10.3390/nul18030392. PMID: 41683216.
 - 120 Zhou J, Qin Y, Xiong X, et al. Effects of iron, vitamin A, and the interaction between the two nutrients on intestinal development and cell differentiation in piglets. *J Anim Sci.* 2021;99:skab258. DOI: 10.1093/jas/skab258. PMID: 34467981.
 - 121 Qin YC, Jin CL, Hu TC, et al. Early Weaning Inhibits Intestinal Stem Cell Expansion to Disrupt the Intestinal Integrity of Duroc Piglets via Regulating the Keap1/Nrf2 Signaling. *Antioxidants (Basel).* 2024;13:1188. DOI: 10.3390/antiox13101188. PMID: 39456442.
 - 122 Kim YH, Jung E, Im GB, et al. Regulation of intracellular transition metal ion level with a pH-sensitive inorganic nanocluster to improve therapeutic angiogenesis by enriching conditioned medium retrieved from human adipose derived stem cells. *Nano Conver.* 2020;7:34. DOI: 10.1186/s40580-020-00244-5. PMID: 33064240.
 - 123 Khalifa YH, Mourad GM, Stephanos WM, et al. Bone Marrow-Derived Mesenchymal Stem Cell Potential Regression of Dysplasia Associating Experimental Liver Fibrosis in Albino Rats. *Biomed Res Int.* 2019;2019:5376165. DOI: 10.1155/2019/5376165. PMID: 31781620.
 - 124 Im GB, Kim YH, Kim YJ, et al. Enhancing the Wound Healing Effect of Conditioned Medium Collected from Mesenchymal Stem Cells with High Passage Number Using Bioreducible Nanoparticles. *Int J Mol Sci.* 2019;20:4835. DOI: 10.3390/ijms20194835. PMID: 31569434.
 - 125 Jalli R, Mehrabani D, Zare S, et al. Cell Proliferation, Viability, Differentiation, and Apoptosis of Iron Oxide Labeled Stem Cells Transfected with Lipofectamine Assessed by MRI. *J Clin Med.* 2023;12:2395. DOI: 10.3390/jcm12062395. PMID: 36983399.
 - 126 Mehrabani D, Nazempour M, Mehdinavaz-Aghdam R, et al. MRI tracking of human Wharton's jelly stem cells seeded onto acellular dermal matrix labeled with superparamagnetic iron oxide nanoparticles in burn wounds. *Burns Trauma.* 2022;10:tkac018. DOI: 10.1093/burnst/tkac018. PMID: 36380853.
 - 127 Iravani K, Mehrabani D, Doostkam A, et al. Use of Magnetic Resonance Imaging to Assess the Regenerative Effects of Adipose Tissue-Derived Mesenchymal Stem Cells in a Rabbit

- Cartilaginous Laryngeal Defect Model. *Curr Ther Res Clin Exp*. 2022;97:100682. DOI: 10.1016/j.curtheres.2022.100682. PMID: 35959231.
- 128 Li X, Wei Z, Li B, et al. In vivo migration of Fe₃O₄@polydopamine nanoparticle-labeled mesenchymal stem cells to burn injury sites and their therapeutic effects in a rat model. *Biomater Sci*. 2019;7:2861-2872. DOI: 10.1039/c9bm00242a. PMID: 31070196.
 - 129 Novotna B, Jendelova P, Kapcalova M, et al. Oxidative damage to biological macromolecules in human bone marrow mesenchymal stromal cells labeled with various types of iron oxide nanoparticles. *Toxicol Lett*. 2012;210:53-63. DOI: 10.1016/j.toxlet.2012.01.008. PMID: 22269213.
 - 130 Li X, Wei Z, Lv H, et al. Iron oxide nanoparticles promote the migration of mesenchymal stem cells to injury sites. *Int J Nanomedicine*. 2019;14:573-589. DOI: 10.2147/IJN.S184920. PMID: 30666115.
 - 131 Zhang C, Liu Z, Zhang Y, et al. "Iron free" zinc oxide nanoparticles with ion-leaking properties disrupt intracellular ROS and iron homeostasis to induce ferroptosis. *Cell Death Dis*. 2020;11:183. DOI: 10.1038/s41419-020-2384-5. PMID: 32170066.
 - 132 Mohammadi S, Palermo A, Ojani P, et al. Comprehensive Effects of Magnesium Supplementation on Cardiometabolic Risk Factors: A Systematic Review and Dose-Response Meta-Analysis. *Nutrients*. 2026;18:2435. DOI: 10.3390/nu18152435. PMID: 42588058.
 - 133 Shimaya M, Muneta T, Ichinose S, et al. Magnesium enhances adherence and cartilage formation of synovial mesenchymal stem cells through integrins. *Osteoarthritis Cartilage*. 2010;18:1300-9. DOI: 10.1016/j.joca.2010.06.005. PMID: 20633668.
 - 134 Kim DH, Shin KK, Jung JS, et al. The Role of Magnesium Ion Substituted Biphasic Calcium Phosphate Spherical Micro-Scaffolds in Osteogenic Differentiation of Human Adipose Tissue-Derived Mesenchymal Stem Cells. *J Nanosci Nanotechnol*. 2015;15:5520-3. DOI: 10.1166/jnn.2015.10463. PMID: 26369111.
 - 135 da Silva Lima F, da Rocha Romero AB, Hastreiter A, et al. An insight into the role of magnesium in the immunomodulatory properties of mesenchymal stem cells. *J Nutr Biochem*. 2018;55:200-208. DOI: 10.1016/j.jnutbio.2018.02.006. PMID: 29554498.
 - 136 Díaz-Tocados JM, Herencia C, Martínez-Moreno JM, et al. Magnesium Chloride promotes Osteogenesis through Notch signaling activation and expansion of Mesenchymal Stem Cells. *Sci Rep*. 2017;7:7839. DOI: 10.1038/s41598-017-08379-y. PMID: 28798480.
 - 137 Ballouze R, Marahat MH, Mohamad S, et al. Biocompatible magnesium-doped biphasic calcium phosphate for bone regeneration. *J Biomed Mater Res B Appl Biomater*. 2021;109:1426-1435. DOI: 10.1002/jbm.b.34802. PMID: 33484103.
 - 138 Yoshizawa S, Brown A, Barchowsky A, et al. Magnesium ion stimulation of bone marrow stromal cells enhances osteogenic activity, simulating the effect of magnesium alloy degradation. *Acta Biomater*. 2014;10:2834-42. DOI: 10.1016/j.actbio.2014.02.002. PMID: 24512978.
 - 139 Rude RK, Gruber HE. Magnesium deficiency and osteoporosis: animal and human observations. *J Nutr Biochem*. 2004;15:710-6. DOI: 10.1016/j.jnutbio.2004.08.001. PMID: 15607643.
 - 140 Castiglioni S, Cazzaniga A, Albisetti W, et al. Magnesium and osteoporosis: current state of knowledge and future research directions. *Nutrients*. 2013;5:3022-33. DOI: 10.3390/nu5083022. PMID: 23912329.
 - 141 Sargenti A, Castiglioni S, Olivi E, et al. Magnesium Deprivation Potentiates Human Mesenchymal Stem Cell Transcriptional Remodeling. *Int J Mol Sci*. 2018;19:1410. DOI: 10.3390/ijms19051410. PMID: 29747379.
 - 142 Loaiza-Giraldo JP, Amigo-Fierro F, Cancino-Castro CI, et al. Effects of Zinc Supplementation on Glycemic Control, Insulin Resistance, Inflammation and Oxidative Stress in Diabetes: A Systematic Review and Meta-Analysis of Randomized Controlled Trials. *Endocrinol Diabetes Metab*. 2026;9:e70264. DOI: 10.1002/edm2.70264. PMID: 42594341.
 - 143 Telgerd MD, Sadeghinia M, Birhanu G, et al. Enhanced osteogenic differentiation of mesenchymal stem cells on metal-organic framework based on copper, zinc, and imidazole coated poly-L-lactic acid nanofiber scaffolds. *J Biomed Mater Res A*. 2019;107:1841-1848. DOI: 10.1002/jbm.a.36707. PMID: 31033136.
 - 144 Lu X, Lin Y, Lin X, et al. Anti-oxidative effect of zinc in human umbilical cord mesenchymal stem cells. *Biophys Rep*. 2021;7:142-151. DOI: 10.52601/bpr.2021.200046. PMID: 37288149.
 - 145 Moon MY, Kim HJ, Choi BY, et al. Zinc Promotes Adipose-Derived Mesenchymal Stem Cell Proliferation and Differentiation towards a Neuronal Fate. *Stem Cells Int*. 2018;2018:5736535. DOI: 10.1155/2018/5736535. PMID: 29765417.
 - 146 Fathi E, Farahzadi R, Rahbarghazi R, et al. Rat

- adipose-derived mesenchymal stem cells aging reduction by zinc sulfate under extremely low frequency electromagnetic field exposure is associated with increased telomerase reverse transcriptase gene expression. *Vet Res Forum*. 2017;8:89-96. PMID: 28785382.
- 147 Xiao L, Chen K, Wen Y, et al. The role of calcium homeostasis dysregulation in allergic rhinitis. *Front Immunol*. 2026;17:1898643. DOI: 10.3389/fimmu.2026.1898643. PMID: 42582083.
 - 148 Gonzalez-McQuire R, Green DW, Partridge KA, et al. Coating of human mesenchymal cells in 3D culture with bioinorganic nanoparticles promotes osteoblastic differentiation and gene transfection. *Adv Mater*. 2007;19:2236-40. DOI: 10.1002/adma.200602770.
 - 149 Hu Q, Tan Z, Liu Y, et al. Effect of crystallinity of calcium phosphate nanoparticles on adhesion, proliferation, and differentiation of bone marrow mesenchymal stem cells. *J Mater Chem*. 2007;17:4690-8. DOI: 10.1039/b710936a.
 - 150 Müller P, Bulnheim U, Diener A, et al. Calcium phosphate surfaces promote osteogenic differentiation of mesenchymal stem cells. *J Cell Mol Med*. 2008;12:281-91. DOI: 10.1111/j.1582-4934.2007.00103.x. PMID: 18366455.
 - 151 Babu D, Ramanathan P, Patel S. Biogenic selenium nanoparticles: a multifunctional tool for combating oxidative stress and chronic diseases. *3 Biotech*. 2026 Aug;16(8):327. doi: 10.1007/s13205-026-04972-z. PMID: 42459412; PMCID: PMC13369280.
 - 152 Rahimi B, Panahi M, Lotfi H, et al. Sodium selenite preserves rBM-MSCs' stemness, differentiation potential, and immunophenotype and protects them against oxidative stress via activation of the Nrf2 signaling pathway. *BMC Complement Med Ther*. 2023;23:131. DOI: 10.1186/s12906-023-03952-7. PMID: 37098557.
 - 153 Liu H, Bian W, Liu S, et al. Selenium protects bone marrow stromal cells against hydrogen peroxide-induced inhibition of osteoblastic differentiation by suppressing oxidative stress and ERK signaling pathway. *Biol Trace Elem Res*. 2012;150:441-50. DOI: 10.1007/s12011-012-9488-4. PMID: 22890880.
 - 154 Ebert R, Ulmer M, Zeck S, et al. Selenium supplementation restores the antioxidative capacity and prevents cell damage in bone marrow stromal cells in vitro. *Stem Cells*. 2006;24:1226-35. DOI: 10.1634/stemcells.2005-0117. PMID: 16424399.
 - 155 Homayoon M, Mehrabani D, Edalatmanesh MA, et al. In Vitro Effect of Lithium Chloride on Adipose Tissue Derived Stem Cells Proliferation and Growth Kinetic. *Iran J Med Sci*. 2026;51:445-454. DOI: 10.30476/ijms.2026.108735.4372. PMID: 42370021.
 - 156 Hedayati A, Homayuon M, Mobaracky A, et al. Lithium Chloride, Ketogenic Diet and Stem Cell Transplantation in Treatment of Bipolar Disorder. *Int J Nutr Sci*. 2024;9:80-82. DOI: 10.30476/IJNS.2024.99601.1250.