



Potential Impact of Probiotics on Low Anterior Resection Syndrome: An Emerging Area of Research

Sara Shojaei-Zarghani¹, PhD;^{ID} Ali Reza Safarpour¹, MD; Alimohammad Banzadeh^{2*}, MD^{ID}

¹Colorectal Research Center, Shiraz University of Medical sciences, Shiraz, Iran

²Laparoscopy Research Center, Shiraz University of Medical Sciences, Shiraz, Iran

***Corresponding author:**

Alimohammad Banzadeh, MD;
Laparoscopy Research Center, Shiraz University of Medical Sciences, Shiraz, Iran.
Email: bananzadeh@gmail.com

Received: 2023-08-27

Revised: 2023-09-13

Accepted: 2023-09-15

Please cite this paper as:

Shojaei-Zarghani S, Safarpour AR, Banzadeh AM. Potential Impact of Probiotics on Low Anterior Resection Syndrome: An Emerging Area of Research. *Iran J Colorectal Res.* 2023;11(3):91-92. doi: 10.30476/ACRR.2023.101187.1201.

The worldwide age-standardized incidence and mortality rates of rectal cancer are estimated to be 7.6 and 3.3 per 100,000, respectively (1). Previous literature suggests that low anterior resection (LAR) may be superior to abdominoperineal resection (APR) for rectal cancer, with better 5-year survival, local recurrence rate, oncological outcomes, and prognosis (2). However, it is reported that around 41% of patients who underwent LAR experience major low anterior resection syndrome (LARS) one year after surgery. LARS is characterized by fecal and gas incontinence, diarrhea, urgency or frequency of stools, sensation of incomplete emptying, and clustering of bowel motions (3). These symptoms are reported to persist over time and influence health related quality of life (4), functional bowel symptoms may also occur after sigmoid resection (5). Despite assessing the effects of several therapeutic strategies in reducing the LARS symptoms, its treatment is still challenging (6).

Perturbation of the gut microbiome has been linked to numerous chronic diseases, such as obesity, endocrine disorders, gastrointestinal diseases, cancer, cardiovascular diseases, etc.(7). Therefore, several research studies have been conducted to assess the effects of modulating the gut microbiome using pre/probiotics. The potential effects of probiotics on colonic disorders complicated by incontinence and abnormal frequency have also been investigated (8, 9). The bidirectional relationship between the gut

microbiome and colonic motility (8) emphasizes the need to assess both the effects of probiotics on LARS and the impact of LARS on gut microbiota.

Few randomized controlled trials (RCT) have assessed the effects of probiotics on LARS score following rectal cancer surgeries with conflicting findings. *Lactobacillus plantarum* CJLP243 as well as VSL#3 (a combination of *Streptococcus thermophilus*, *Bifidobacterium breve*, *Bifidobacterium longum*, *Bifidobacterium infantis*, *Lactobacillus acidophilus*, *Lactobacillus casei*, *Lactobacillus delbrueckii* subsp. *bulgaricus* and *Lactobacillus plantarum*) supplementation had no effects on bowel motions and LARS score when administered in the perioperative period of ileostomy reversal or exactly following reversal of a temporary loop ileostomy (10, 11). However, in another RCT, probiotic administration (*Bifidobacterium animalis* subsp. *lactis* HY8002, *Lactobacillus casei* HY2782, and *Lactobacillus plantarum* HY7712), starting one week before surgery and continuing for 21 days after anterior sigmoid colon cancer resection, improved flatus control compared to the placebo. Nonetheless, it had no effect on the total LARS score (12). Differences in the site of cancer, type and dosage of probiotics, and timing following surgery or stoma reversal could explain this conflicting evidence.

A recent study evaluated gut microbiome 13.3 and 2.2 months after LAR and ileostomy closure, respectively. A decreased diversity of gut microbiome,

a higher level of *Bacteroidaceae* enterotype, and lower levels of *Prevotellaceae* enterotype and lactic acid-producing bacteria, including *Lactobacillus* and *Bifidobacterium*, were detected in patients with frequency-dominant LARS. The authors suggested that modulating gut microbiota could be a potential therapeutic option for patients experiencing symptoms related to frequency, including frequency, clustering, and urgency, but not for those with an incontinence-dominant pattern of LARS (13).

Based on the existing literature, it is recommended that the long-term effects of pre/probiotics be assessed after overcoming the acute phase of LARS, which typically occurs immediately following surgery and stoma reversal. Research should also investigate the effects of probiotics on various symptoms of LARS and their impact on the gut microbiome.

Conflict of interest: None declared.

References

1. Global Cancer Observatory: Cancer Today. Lyon, France: International Agency for Research on Cancer. Available from: <https://gco.iarc.fr/today/data/factsheets/cancers/9-Rectum-fact-sheet.pdf>.
2. Wang XT, Li DG, Li L, Kong FB, Pang LM, Mai W. Meta-analysis of oncological outcome after abdominoperineal resection or low anterior resection for lower rectal cancer. *Pathology oncology research : POR*. 2015;21(1):19-27.
3. Croese AD, Lonie JM, Trollope AF, Vangaveti VN, Ho YH. A meta-analysis of the prevalence of Low Anterior Resection Syndrome and systematic review of risk factors. *International journal of surgery (London, England)*. 2018;56:234-41.
4. Pieniowski EHA, Palmer GJ, Juul T, Lagergren P, Johar A, Emmertsen KJ, et al. Low Anterior Resection Syndrome and Quality of Life After Sphincter-Sparing Rectal Cancer Surgery: A Long-term Longitudinal Follow-up. *Diseases of the colon and rectum*. 2019;62(1):14-20.
5. Bryant CLC, Lunniss PJ, Knowles CH, Thaha MA, Chan CLH. Anterior resection syndrome. *Lancet Oncol*. 2012;13(9):e403-8.
6. Emile SH, Garoufalia Z, Barsom S, Horeh N, Gefen R, Zhou P, et al. Systematic review and meta-analysis of randomized clinical trials on the treatment of low anterior resection syndrome. *Surgery*. 2023;173(6):1352-8.
7. Afzaal M, Saeed F, Shah YA, Hussain M, Rabail R, Socol CT, et al. Human gut microbiota in health and disease: Unveiling the relationship. 2022;13.
8. Ma W, Drew DA, Staller K. The Gut Microbiome and Colonic Motility Disorders: A Practical Framework for the Gastroenterologist. *Current gastroenterology reports*. 2022;24(10):115-26.
9. Liu L, Wang A, Shi H, Tao H, Nahata MC. Efficacy and safety of probiotics and synbiotics for functional constipation in children: A systematic review and meta-analysis of randomized clinical trials. *Clinical nutrition (Edinburgh, Scotland)*. 2023;42(10):1817-26.
10. Stephens JH, Hewett PJ. Clinical trial assessing VSL#3 for the treatment of anterior resection syndrome. *ANZ journal of surgery*. 2012;82(6):420-7.
11. Yoon BJ, Oh HK, Lee J, Cho JR, Kim MJ, Kim DW, et al. Effects of probiotics on bowel function restoration following ileostomy closure in rectal cancer patients: a randomized controlled trial. *Colorectal disease : the official journal of the Association of Coloproctology of Great Britain and Ireland*. 2021;23(4):901-10.
12. Park IJ, Lee J-H, Kye B-H, Oh H-K, Cho YB, Kim Y-T, et al. Effects of probiotics on the symptoms and surgical outcomes after anterior resection of colon cancer (POSTCARE): a randomized, double-blind, placebo-controlled trial. *Journal of clinical medicine*. 2020;9(7):2181.
13. Kim MJ, Park S, Park JW, Choi J, Kim HJ, Lim HK, et al. Gut microbiome associated with low anterior resection syndrome after rectal cancer surgery. *Scientific reports*. 2023;13(1):8578.